Serbian Biochemical Society

President: Marija Gavrović-Jankulović **Vice-president:** Suzana Jovanović-Šanta **General Secretary:** Isidora Protić-Rosić **Treasurer:** Milica Popović

Scientific Board

Marija Gavrović-Jankulović Svetlana Dinić Ario de Marco Suzana Jovanović-Šanta Mario Gabričević Vladimir Mihailović Theodore G. Sotiroudis Natalija Polović Andreja Rajković Nataša Simin Edvard Petri Sanja Krstić Željko Popović Snežana Pantović Milan Nikolić Simeon Minić

Organization Committee

Ivan Spasojević Tanja Ćirković Veličković Milica Popović Aleksandra Uskoković Tijana Ćulafić Isidora Protić-Rosić Jovana Trbojević-Ivić Milena Dimitrijević Srđan Miletić

Proceedings

Editor: Ivan Spasojević Technical support: Jovana Trbojević-Ivić, Milena Dimitrijević, Tijana Ćulafić Cover design: Zoran Beloševac Publisher: Faculty of Chemistry, Serbian Biochemical Society Printed by: Colorgrafx, Belgrade No of printed copies: 130

Serbian Biochemical Society Twelfth Conference

International scientific meeting

September 21-23, 2023, Belgrade, Serbia

"Biochemistry in Biotechnology"

Applying immobilised phycobiliproteins onto chitosan for efficient mercury removal

Jelena Radović¹, Dragana Popović², Tatjana Ćurčić², Milan Nikolić^{1,3}, Simeon Minić^{1,3}, Nikola Gligorijević^{4*}

¹Department of Biochemistry, Faculty of Chemistry, University of Belgrade, Serbia ²Institute for Biocides and Medical Ecology, Belgrade, Serbia ³Center of Excellence for Molecular Food Sciences, Faculty of Chemistry, University of Belgrade ⁴Department of Chemistry, Institute of Chemistry, Technology, and Metallurgy, National

Institute of the Republic of Serbia, University of Belgrade

*e-mail: nikola.gligorijevic@ihtm.bg.ac.rs

This study aimed to improve chitosan polymer's capabilities to absorb mercury by immobilising phycobiliproteins (PBPs) onto the surface of chitosan beads (chitosan-PBPs). Phycobiliproteins, light-harvesting proteins from algae and cyanobacteria, have several industrially essential applications. These proteins can bind heavy metals with high affinities. Protein extracts obtained from both Arthrospira platensis, with C-phycocyanin as the dominant phycobiliprotein and Neoporphyra haitanensis, with R-phycocyanin and R-phycoerythrin as the dominant PBPs, were covalently immobilised onto chitosan beads. Binding analysis showed that, on average, 54 µg of PBPs were immobilised per bead. Immobilised proteins were still in their native state, with no visible colour change after immobilisation. Chitosan-PBPs and chitosan alone were tested for mercury adsorption at pH 4 and pH 7 by atomic absorption spectroscopy. The tested concentration range of mercury was from 1 to 70 ppm. Affinity, calculated using Henry's binding isotherm, of chitosan-PBPs for mercury was higher at both pH values than chitosan alone. Furthermore, chitosan–PBPs beads were able to absorb significantly more mercury than chitosan alone. These results show that the covalent immobilisation of PBPs onto chitosan improves its mercury adsorption characteristics and creates a more efficient eco-friendly adsorbent for removing mercury ions in the tested concentration range.

Acknowledgements

This study was financially supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia, contract numbers: 451-03-47/2023-01/200168 and 451-03-47/2023-01/200026.

CIР - Каталогизација у публикацији Народна библиотека Србије, Београд

577.1(048)

SERBIAN Biochemical Society. International scientific meeting (12; 2023; Beograd)

"Biochemistry in Biotechnology" : [proceedings] / Serbian Biochemical Society, Twelfth Conference, International scientific meeting, September 21-23, 2023, Belgrade, Serbia ; [editor Ivan Spasojević]. - Belgrade : Faculty of Chemistry : Serbian Biochemical Society, 2023 (Belgrade : Colorgrafx). - 156 str. ; 23 cm

Tiraž 130. - Bibliografija uz većinu apstrakata.

ISBN 978-86-7220-140-6 (FOC)

а) Биохемија -- Апстракти

COBISS.SR-ID 124201993
