10th IAPC Meeting

Tenth World Conference on Physico-Chemical Methods in Drug Discovery &

Sixth World Conference on ADMET and DMPK

Book of Abstracts



10th IAPC Meeting

Tenth World Conference on Physico-Chemical Methods in Drug Discovery & Sixth World Conference on ADMET and DMPK Belgrade, Serbia, September 4-6, 2023

Book of Abstracts

Organized by International Association of Physical Chemists & Faculty of Chemistry, University of Belgrade, Serbia

Published by International Association of Physical Chemists E-mail: office@iapchem.org, URL: http://www.iapchem.org

For Publisher **Zoran Mandić**

Editor **Tatjana Verbić & Zoran Mandić**

Design, page making and computer layout Aleksandar Dekanski

On Line version only

The Scientific and Organizing Committee:

Tatjana Verbić, Conference Chair University of Belgrade, Serbia,

Alex Avdeef ADME Research, New York, USA

Kiyohiko Sugano Ritsumeikan University, Osaka, Japan

> Kin Tam University of Macau, Macau

Zoran Mandić University of Zagreb, Croatia

Klara Valko Biomimetic chromatography Ltd. UK

Godefridus J. Peters Amsterdam University Medical Centers, The Netherlands

> Hong Wan WHDeX Consultng AB, Sweden

Local Organizing Committee

Tatjana Verbić, Conference Chair University of Belgrade, Faculty of Chemistry, Serbia

Goran Roglić University of Belgrade, Faculty of Chemistry, Serbia

Ilija Cvijetić University of Belgrade, Faculty of Chemistry, Serbia

Miloš Pešić University of Belgrade, Faculty of Chemistry, Serbia

Olivera Marković University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

Aleksandar Dekanski University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

> Marija Popović Nikolić University of Belgrade, Faculty of Pharmacy, Serbia

Organization of IAPC-10 Meeting is supported by



Ministry of Science, Technological Development and Innovation of the Republic of Serbia

Sponsors and Exhibitors





<u>N</u>enad M. Kostić Fund for Chemical Sciences



























¹H-NMR metabonomic view on schizophrenia

Nataša Avramović^{1,*}, Katarina Simić², Zoran Miladinović³, Nina Todorović², Snežana Trifunović⁴, Aleksandra Gavrilović⁵, Silvana Jovanović⁵, Dejan Gođevac², Ljubodrag Vujisić⁴, Vele Tešević⁴, Ljubica Tasić⁶, Boris Mandić⁴

 ¹University of Belgrade-Faculty of Medicine, Institute of Medical Chemistry, Višegradska 26, 11000 Belgrade, Serbia
²Institute of Chemistry, Technology and Metallurgy, National Institute, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia
³Institute of General and Physical Chemistry, Studentski trg 12-16, 11158 Belgrade, Serbia;
⁴University of Belgrade - Faculty of Chemistry, Studentski trg 12-16, 11000 Belgrade, Serbia
⁵Special Hospital for Psychiatric Diseases "Kovin", Cara Lazara 253, 26220 Kovin, Serbia
⁶ Institute of Chemistry, Organic Chemistry Department, State University of Campinas, Campinas 13083-970, SP, Brazil
*Correspondence: natasa.avramovic@med.bg.ac.rs

Schizophrenia (SCZ) is a brain disease leading to significant functional impairments and premature death, and it affects 20 million people worldwide. Due to the complexity of this disease including different genetic and environmental factors, there is a lack in understanding pathophysiology and diagnosis of schizophrenia. In order to overcome existing gaps, the establishment of a universal set of SCZ biomarkers has a crucial role. Metabonomic study of serum samples of Serbian patients with schizophrenia (51) and healthy controls (39) by ¹H-NMR analyses associated with chemometrics, provided the identification of 26 metabolites/biomarkers for this disorder. The biomarker set including aspartate/aspartic acid, lysine, 2-hydroxybutyric acid, and acylglycerols was established for the first-time in SCZ serum samples of Serbian patients by ¹H-NMR experiments. The other 22 identified metabolites are in agreement with the previously confirmed NMR-based serum biomarker sets of Brazilian and/or Chinese patient samples. The same 13 metabolites (lactate/lactic acid, threonine, leucine, isoleucine, valine, glutamine, asparagine, alanine, gamma-aminobutyric acid, choline, glucose, glycine and tyrosine) were established in all SCZ samples from three countries of different ethnicity and geographical origins (Serbia, Brazil and China). These results emphasize the crucial role in the possibility of their application as biomarkers for diagnosis of SCZ, reliable monitoring of treatment response and clinical outcomes.