

## SUPPLEMENTARY MATERIAL

# New aurone epoxide and auronolignan from the heartwood of *Cotinus coggygria* Scop.

Miroslav Novakovic<sup>1\*</sup>, Iris Djordjevic<sup>2</sup>, Nina Todorovic<sup>1</sup>, Snezana Trifunovic<sup>3</sup>, Boban Andjelkovic<sup>3</sup>, Boris Mandic<sup>3</sup>, Milka Jadranin<sup>1</sup>, Ivan Vuckovic<sup>4</sup>, Vlatka Vajs<sup>1</sup>, Slobodan Milosavljevic<sup>3</sup> and Vele Tesevic<sup>3</sup>

<sup>1</sup>*Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia (e-mail: [mironov76@yahoo.com](mailto:mironov76@yahoo.com); [ninat@chem.bg.ac.rs](mailto:ninat@chem.bg.ac.rs); [milkaj@chem.bg.ac.rs](mailto:milkaj@chem.bg.ac.rs); [vvajs@chem.bg.ac.rs](mailto:vvajs@chem.bg.ac.rs))*

<sup>2</sup>*Faculty of Veterinary Medicine, University of Belgrade, Bulevar oslobođenja 18, 11000 Belgrade, Serbia (e-mail: [iris@vet.bg.ac.rs](mailto:iris@vet.bg.ac.rs))*

<sup>3</sup>*Faculty of Chemistry, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia ([snezanan@chem.bg.ac.rs](mailto:snezanan@chem.bg.ac.rs); [aboban@chem.bg.ac.rs](mailto:aboban@chem.bg.ac.rs); [borism@chem.bg.ac.rs](mailto:borism@chem.bg.ac.rs); [smilo@chem.bg.ac.rs](mailto:smilo@chem.bg.ac.rs); [vtesevic@chem.bg.ac.rs](mailto:vtesevic@chem.bg.ac.rs))*

<sup>4</sup>*Mayo Clinic, College of Medicine, Rochester, MN 55905, USA ([vuckovic.ivan@mayo.edu](mailto:vuckovic.ivan@mayo.edu))*

\*Corresponding author, e-mail: [mironov76@yahoo.com](mailto:mironov76@yahoo.com)

\*Corresponding author, e-mail: [mironov76@yahoo.com](mailto:mironov76@yahoo.com)

**Abstract.** From the methylene chloride/methanol extract of *Cotinus coggygria* Scop. heartwood one new aurone epoxide, 2,10-oxy-10-methoxysulfuretin (**14**), and one auronolignan (**15**), named cotinignan A, were isolated by silica gel column and semipreparative HPLC chromatography. In addition, thirteen known secondary metabolites namely sulfuretin, 2,3-*trans*-fustin, fisetin, butin, butein, taxifolin, eriodictyol, 3',5,5',7-tetrahydroxyflavanone, 3',4',7-trihydroxyflavone, 3-*O*-methyl-2,3-*trans*-fustin, 3-*O*-galloyl-

*2,3-trans*-fustin,  $\beta$ -resorcylic acid and 3-*O*- $\beta$ -sitosterol glucoside were isolated as well. Their structures were elucidated by 1D and 2D NMR, HR-ESI-MS, IR and UV. Ten out of eleven isolated flavonoids possess 7, 3' and 4' hydroxy groups. These structural features could be considered as chemotaxonomic characteristic of flavonoids from *C. coggygria*. Cotinignan A (**15**) represents new subclass of secondary metabolites - auronolignans.

**Keywords.** *Cotinus coggygria* Scop, flavonoids, cotinignan A, semipreparative HPLC, NMR

**Table S1.** HPLC program for the quantification of the main compounds from the *C. coggygria* CH<sub>2</sub>Cl<sub>2</sub>/MeOH extract

**Figure S1.** UV spectra of **14** in MeOH, with addition of AlCl<sub>3</sub>, and HCl

**Figure S2.** UV spectra of compound **14** in MeOH, with NaOAc, NaOAc+H<sub>3</sub>BO<sub>3</sub> and with NaOMe

**Figure S3.** <sup>1</sup>H NMR spectrum of compound **14**

**Figure S4.** <sup>13</sup>C NMR spectrum of compound **14**

**Figure S5.** Aromatic part of the COSY spectrum of **14**

**Figure S6.** Aromatic part of the NOESY spectrum of **14**

**Figure S7.** Aromatic part of the HSQC spectrum of **14**

**Figure S8.** Aromatic part of the HMBC spectrum of **14**

**Figure S9.** Key HMBC correlations of **14**

**Table S2.** <sup>1</sup>H, <sup>13</sup>C NMR and HMBC data of **14** (CD<sub>3</sub>OD,  $\delta$  ppm, *J* in Hz)

**Figure S10.** Aromatic part of the <sup>1</sup>H NMR spectrum of **15**

**Figure S11.** Aliphatic part of the <sup>1</sup>H NMR spectrum of **15**

**Figure S12.** <sup>13</sup>C NMR spectrum of **15**

**Figure S13.** HSQC spectrum of **15**

**Figure S14.** The first part of the HMBC spectrum of **15**

**Figure S15.** The second part of the HMBC spectrum of **15**

**Figure S16.** HMBC (4 Hz) correlation H-7''/C-4' (**15**)

**Figure S17.** The first part of the NOESY spectrum of **15**

**Figure S18.** The second part of the NOESY spectrum of **15**

**Table S3.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of **15** ( $\text{CD}_3\text{OD}$ ,  $\delta$  ppm,  $J$  in Hz)

**Figure S19.** 3-Aryl-propanol moiety in **15**


**Figure S20.** Key HMBC correlations in **15**

**Figure S21.** UV spectrum of **15**


**Figure S22.** UV spectrum of **10**

**Table S1.** HPLC program for the quantification of the main compounds from the *C. coggygria*  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  extract


t (min)	A (water)	B (acetonitrile)	Flow (mL/min)
0	95	5	1.4
1.5	95	5	1.4
26	5	95	1.4
35	5	95	1.4
36	95	5	1.4
41	95	5	1.4




**Figure S1.** UV spectra of **14** in MeOH, with addition of AlCl<sub>3</sub>, and HCl




**Figure S2.** UV spectra of **14** in MeOH, with NaOAc, NaOAc+H<sub>3</sub>BO<sub>3</sub> and with NaOMe




**Figure S3.**  $^1\text{H}$  NMR spectrum of **14**




**Figure S4.**  $^{13}\text{C}$  NMR spectrum of **14**




**Figure S5.** Aromatic part of the COSY spectrum of **14**




**Figure S6.** Aromatic part of the NOESY spectrum of **14**



**Figure S7.** Aromatic part of the HSQC spectrum of **14**




**Figure S8.** Aromatic part of the HMBC spectrum of **14**




**Figure S9.** Key HMBC correlations of **14**

**Table S2.**  $^1\text{H}$ ,  $^{13}\text{C}$  NMR and HMBC data of **14** ( $\text{CD}_3\text{OD}$ ,  $\delta$  ppm,  $J$  in Hz)


Position	$\delta_{\text{H}}$	$\delta_{\text{C}}$	HMBC $_{\text{H}}\rightarrow_{\text{C}}$
2		90.97	
3		190.16	
4	7.67 d (8.8 Hz)	128.88	8, 6
5	6.48 dd (8.8; 2.2 Hz)	110.91	7
6		165.40	
7	6.35 d (2.2 Hz)	103.37	9
8		158.74	
9		111.87	
10		107.26	
1'		124.66	
2'	7.17 d (2.0 Hz)	116.99	10, 4', 6'
3'		144.34	
4'		146.02	
5'	6.79 d (8.4 Hz)	114.61	1', 3'
6'	7.04 dd (8.4; 2.0 Hz)	120.51	2', 4'
10-OMe	3.03 s	50.04	10




**Figure S10.** Aromatic part of the  $^1\text{H}$  NMR spectrum of **15**




**Figure S11.** Aliphatic part of the  $^1\text{H}$  NMR spectrum of **15**




**Figure S12.**  $^{13}\text{C}$  NMR spectrum of **15**




**Fig. S13.** HSQC spectrum of **15**




**Figure S14.** The first part of the HMBC spectrum of **15**




**Figure S15.** The second part of the HMBC spectrum of **15**



**Figure S16.** HMBC (4 Hz) correlation H-7''/C-4' (**15**)



**Figure S17.** The first part of the NOESY spectrum of **15**




**Figure S18.** The second part of the NOESY spectrum of **15**


**Table S3.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of compounds **15** ( $\text{CD}_3\text{OD}$ ,  $\delta$  ppm,  $J$  in Hz)

Compound	<b>15</b>	
Position	$\delta_{\text{H}}$	$\delta_{\text{C}}$
2	-	148.4
3	-	184.5
4	7.60 d (8.5)	128.2
5	6.68 dd (8.5; 2.0)	114.4
6	-	168.9
7	6.70 d (2.0)	99.6
8	-	170.1
9	-	114.6
10	6.71 s	113.4
1'	-	127.0
2'	7.62 d (2.0)	120.8
3'	-	145.5
4'	-	147.0
5'	7.04 d (8.5)	118.5
6'	7.39 dd (8.5; 2.0)	127.0


1"	-	128.3
2"	6.76 s	106.0
3"	-	149.5
4"	-	137.3
5"	-	149.5
6"	6.76 s	106.0
7"	4.95 d (8.0)	78.0
8"	4.15 ddd (8.0; 4.0; 2.5)	80.4
9"	3.75 dd (12.5; 2.5) 3.51 dd (12.5; 4.0)	62.0
3"-OMe	3.89 s	57.0
5"-OMe	3.89 s	57.0




**Figure S19.** 3-Aryl-propanol moiety in **15**



**Figure S20.** Key HMBC correlations in **15**



**Figure S21.** UV spectrum of **15**



**Figure S22.** UV spectrum of **10**