

EMEC21

21st European Meeting on Environmental Chemistry
November 30 – December 3, 2021, Novi Sad, Serbia

www.emec21.rs



Association of Chemistry
and the Environment



Serbian Chemical Society



Matica Srpska

Scientific Committee

Jan Schwarzbauer, president

Organisational Committee


Branimir Jovančević, president

Executive Committee

Vladimir Beškoski, president



BOOK OF ABSTRACTS



21st European Meeting
on Environmental Chemistry

EMEC 21



21st European Meeting
on Environmental Chemistry

BOOK OF ABSTRACTS
EMEC 21

November 30 – December 3, 2021

Novi Sad, Serbia



Book of Abstracts
21st European Meeting on Environmental Chemistry

Publisher

Serbian Chemical Society
Karnegijeva 4/III, Belgrade, Republic of Serbia

For the publisher

Dušan Sladić
President of the Serbian Chemical Society

Editors

Ivana Ivančev-Tumbas
Vladimir P. Beškoski
Aleksandra Šajnović

Cover page photo

Branko Lučić

Design and prepress

Beoživković, Belgrade

Printed by

RIS Studio, Belgrade

Circulation

150

ISBN

978-86-7132-078-8

Year

2021

Scientific Committee

Jan Schwarzbauer, Germany, president

Ivana Ivančev-Tumbas, Serbia, vice president

Marijana Ačanski, Serbia	Đorđe Jovanović, Serbia
Vladimir P. Beškoski, Serbia	Albert T. Lebedev, Russia
Jelena Bošković, Serbia	Milan Matavulj, Serbia
Anne-Marie Delort, France	Milica Balaban, Bosnia and Herzegovina
Stuart Gibb, UK	Polonca Trebše, Slovenia
Branimir Jovančićević, Serbia	

Organisational Committee

Branimir Jovančićević, president

Maja Turk Sekulić, vice president

Mira Aničić-Urošević	Jovana Orlić
Mališa Antić	Kristijan Pastor
Jelena Avdalović	Ivana Perović
Tanja Brdarić	Petar Pfindt
Aleksandar Đorđević	Srđan Pokorni
Rada Đurović Pejčev	Bojan Radak
Gordana Gajica	Dubravka Relić
Ljubiša Ignjatović	Goran Roglić
Marko Ilić	Sanja Sakan
Konstantin Ilijević	Sladana Savić
Ivana Ivančev-Tumbas	Jelena Savović
Kristina Joksimović	Slavka Stanković
Vladan Joldžić	Vesna Stanković
Milica Kašanin-Grubin	Sanja Stojadinović
Branka Lončarević	Aleksandra Šajnović
Nikoleta Lugonja	Tatjana Šolević Knudsen
Marija Lješević	Anđelka Tomašević
Snežana Maletić	Aleksandra Tubić
Aleksandra J. Mihajlidi-Zelić	Gorica Veselinović
Maja Milanović	Nenad Zarić
Srđan Miletić	Vesna Zlatanović Tomašević
Tijana Miličević	Aleksandra Žerađanin
Dubravka Milovanović	Sanja Živković
Miloš Momčilović	

Executive Committee

Vladimir P. Beškoski, president

Jelena Radonić, vice president

Gordana Gajica

Kristina Joksimović

Milica Kašanin-Grubin

Srđan Miletić

Gorica Veselinović

EMEC21 Secretariat

Aleksandra Šajnović

Aleksandra Tubić

Nevena Antić

Soil Contamination by Petroleum Products in the Urban Environments: Belgrade, Serbia

G. Devic¹, M. Ilic¹, S. Bulatovic², S. Miletic¹, M. M. Vrvic². (1) University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11000 Belgrade, Serbia, (2) Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia; gordana.devic.@ihtm.bg.ac.rs and gdevic@chem.bg.ac.rs.

Environmental pollution with petroleum and petrochemical products has attracted much attention in recent years. The present study is focused on the investigation of urban soil pollution in the area of thermal plant New Belgrade with more than 200000 residents. Thermal Power Plant New Belgrade is located on the left bank of the Sava, about 1 km from the Sava's confluence with the Danube. The Thermal Power Plant complex consists of storage tanks for crude oil and oil products and it was contaminated due to break-down of the mazut reservoirs (2009, during a gas crisis) and NATO bombing of the reservoirs (1999).

A total of 45 soil samples were collected in May, 2015, and in total 8 geochemical parameters were determined by using official or recommended methods [1].

Statistical data processing includes application of multivariate statistical methods to previously systematized data on geochemical parameters in soil samples from Belgrade city. Two multivariate statistical methods have been applied – hierarchical cluster analysis (HCA) and factor analysis – analysis of the main components (PCA).

From the output of the hierarchical cluster analysis, a total of three clusters of the soil samples were recognized according to the level of clustering. Cluster A and B are linked at a shorter distance and are together linked to Cluster C at a longer distance. Component 1 of the PCA dominated in group A. Samples of group A contained the maximum amounts of total aliphatic hydrocarbon (TPH) (71.85 ± 17.9), and Unresolved/Resolved complex mixture (U/R) (3.02 ± 1.8) higher than the other groups. The results of the soil analyses indicated that most samples classified as C1 may reflect anthropogenic contamination of the urban environment. The results showed that the top soil contained high concentration of TPH (90.65 mg kg^{-1}). This may be a result was caused by crude oil spills and leaks which continuously occurred for long periods of time. According to the "Regulation on the program of systematic monitoring of land quality, indicators for According to the "Regulation on the program of systematic monitoring of land quality, indicators for assessment of the risk of land degradation and the methodology for making remediation programs "(Official Gazette of RS, No. 88/2010-Annex 13) are not higher than" values that may indicate significant

contamination "–remediation values, but are higher than "limit values", which means that this area must be programs "(Official Gazette of RS, No. 88/2010-Annex 13) are not higher than" values that may indicate significant contamination "–remediation values, but are higher than" limit values ", which means that this area must be under permanently monitoring.

Group B is represented by a principal component 2 related to the highest CPI (3.48 ± 1.32), Low/High alkanes (0.299 ± 0.45), and ACL (the average number of carbon atoms per molecule based on the abundance of odd-carbon-numbered higher plant *n*-alkanes) (29.67 ± 0.40). The analyzed samples have similar ACL contents in all groups. Distribution of these parameters suggesting predominant biogenic sources rather than petroleum related input.

The component 3 of the PCA2, loaded by *n*-C17/ Pr, and *n*-C18/Phy, also constituted a strong cluster C, *n*-C17 and *n*-C18 are more abundant than the isoprenoids pristane and phytane. Comparatively, isoprenoid hydrocarbons are more resistant to biodegradation than *n*-alkanes [2], leading to the increase of Pr/ *n*-C17 and Ph/*n*-C18 ratios to a value of much higher than 1 when they have been deeply degraded [3]. The distribution of Pr/*n*-C17, and Ph/*n*-C18 ratios for almost all samples are small than 1, indicate biogenic contribution, except of some samples are around 1, points to petroleum contamination source in urban area of Belgrade city.

Acknowledgements

The authors would like to thank the Ministry of Education, Science and Technological Development (Grant Nos. III 43004).

References

- [1] B, Jovančićević, M, Antić., T. Šolević, MM. Vrvic, A. Kronimus, J. Schwarzbauer. Environmental Science and Pollution Research, 12, (2005), 205–212.
- [2] S.J. Varjani, Bioresour. Technol. (2017), 223, 277–286.
- [3] F.J. Gonza' lez-Vila, O. Polvillo, T. Boski, J.R Moura, D., Andre' s de, Organic Geochemistry 34, (2003), 1601–1613.