Serbian Biochemical Society

President: Marija Gavrović-Jankulović Vice-president: Suzana Jovanović-Šanta General Secretary: Isidora Protić-Rosić

Treasurer: Milica Popović

Scientific Board

Marija Gavrović-Jankulović

Mihajlo B. Spasić Vesna Niketić

Ivanka Karadžić Svetlana Dinić

Nevena Đukić Jelena Bašić Ivan Spasojević

Ivana Beara Mojca Stojiljković Andjelka Čelić

Željko Popović

Žanka Bojić Trbojević

Milan Nikolić

Ana Ninić

Adela Pitea Zupkó István Vlatka Zoldos Aleksandra Inić-

Kanada

Tomasz Jurkowski Yaraslau Dzichenka Brankica Janković Sanja Krstić

Organization Committee

Suzana Jovanović-

Šanta

Jelena Purać

Milica Popović Emilija Svirčev

Miloš Opačić Milena Dimitrijević

Tatjana Majkić Sofija Bekić Diandra Pintać

Isidora Protić-Rosić Marina Crnković Maja Marinović

Iva Uzelac Jovana Drljača Miloš Avramov

Srđana Đorđievski Milana Bosanac

Vanja Tatić

Proceedings

Editor: Ivan Spasojević

Technical support: Jelena Korać Jačić

Cover design: Zoran Beloševac

Publisher: Faculty of Chemistry, Serbian Biochemical Society

Printed by: Colorgrafx, Belgrade

Serbian Biochemical Society Eleventh Conference

Scientific meeting of an international character

September 22nd and 23rd, 2022, Novi Sad, Serbia

"Amazing Biochemistry"

Production and application of pectinases in the liquefaction of apricot and blueberry juice

Marija Pavlović^{1,2*}, Aleksandra Margetić³, Marinela Šokarda Slavić³, Marina Ristović³, Ratko Pavlović², Stefan Nikolić⁴, Zoran Vujčić²

Pectinases are widely used in the fruit juice industry for clarification, liquefaction and stabilization of juices¹. One of the biggest problems in the production of fruit juices is the turbidity of the juice, which is mainly caused by the presence of pectin polysaccharides. Therefore, pectinase is used in juice clarification, which breaks down the pectin structure and reduces unwanted cloudiness and sediment². In this work, the production of pectinases was optimized by solid state fermentation using *Aspergillus tubingensis* strain, which proved to be an efficient producer of these enzymes. Statistical method Design of Experiment was used to optimize the medium and conditions for enzyme production. The total pectinase activity obtained was determined by the DNS method (47 U/mL). Endopectinases activity is determined by reduction of viscosity of pectin solutions. The resulting complex of pectinase enzymes was used for the liquefaction of apricot and blueberry pulp, with a juice yield of 72% and 81%, respectively. Also, apricot juice treated with enzymes was clarified by 77% compared to juice that was not treated with enzymes. Blueberry juice obtained after treatment with pectinase enzymes has a higher antioxidant activity than the untreated juice, as determined by the DPPH assay.

Acknowledgements

This work was financially supported by The Ministry of Education, Science and Technological Development of the Republic of Serbia, Contract numbers: 451-03-68/2022-14/200168, 451-03-68/2022-14/200026, 451-03-9/2022-14/200017 and 451-03-68/2022-14/200288.

References

 Sandri IG, Lorenzoni CMT, Fontana RC, Silveira MM. Use of pectinases produced by a new strain of Aspergillus niger for the enzymatic treatment of apple and blueberry juice. LWT -Food Sci Technol 2013;51:469-75.

¹Department of Physical Chemistry, Vinča Institute of Nuclear Sciences, National Institute of the Republic of Serbia, University of Belgrade, Belgrade, Serbia

²Department of Biochemistry, Faculty of Chemistry, University of Belgrade

³Department of Chemistry, Institute of Chemistry, Technology and Metallurgy, National Institute of the Republic of Serbia, University of Belgrade

⁴Innovative Centre of the Faculty of Chemistry, University of Belgrade

^{*}e-mail: marija.pavlovic@vin.bg.ac.rs

2.	Mahmoodi M, Najafpour GD, Mohammadi M. Production of pectinases for quality apple juice through fermentation of orange pomace. J Food Sci Technol 2017;54;4123-8.