

PRELIMINARY COMMUNICATION

**Synthesis of a steroidal dendrimer core**

TATJANA KOP<sup>1#</sup>, GABRIELLA POCSFALVI<sup>2</sup> and BOGDAN A. ŠOLAJA<sup>1#\*</sup>

<sup>1</sup>*Faculty of Chemistry, University of Belgrade, P. O. Box 158, 11001 Belgrade, Serbia and Montenegro (E-mail: bsolaja@chem.bg.ac.yu) and* <sup>2</sup>*Istituto di Scienze dell'Alimentazione, Consiglio Nazionale delle Ricerche, Avellino, Italy*

(Received 26 February 2004)

*Abstract:* Synthesis of a steroidal dendrimer core possessing various functional termini, such as ester, carboxy and hydroxy, is presented. The approach described enables further simple manipulations for the introduction of more complex functionalities.

*Keywords:* deoxycholic acid, trimesoyl trichloride, core, dendrimer.

INTRODUCTION

Dendrimers are well-defined macromolecules of uniform mass that contain a core, successive layers of branched repeat units and surface groups.<sup>1</sup> Over the last decade or more, the synthesis of novel dendrimers has been a very active area of research and one important aspect has been the incorporation of functional units such as crown ethers,<sup>2</sup> mesogenic groups<sup>3</sup> and various redox-active substituents based on tetrathiafulvalene,<sup>4</sup> anthraquinones,<sup>5</sup> ferrocene,<sup>6</sup> fullerenes,<sup>7</sup> or transition metal complexes.<sup>8</sup> An interesting consequence of placing a functional unit at the core of a dendrimer is its steric isolation which can prevent unwanted interference of the functionality.<sup>9</sup>

The three-dimensional globular morphology of dendrimer macromolecules<sup>1</sup> makes them attractive for higher level structural ordering in a manner resembling the folded structures of proteins.<sup>2</sup> Many of the potential applications envisaged for dendrimers (*e.g.*, enantioselective catalysts, chemical sensors, optical switches) can only be realized in materials expressing highly ordered and controllable conformations. However, the goal of inducing higher-order three-dimensional organization in chiral dendrimers<sup>3</sup> has been severely hampered by the conformational flexibility of most commonly studied dendrimers.<sup>4</sup>

Recent results of our research in antimalarial peroxide chemistry<sup>10</sup> prompted us to envisage steroidal-based dendrimers which would act as a carrier of the tetraoxane function-

# Serbian Chemical Society active member.


\* corresponding author: Phone: +381-11-63-86-06. Fax: +381-11-63-60-61

ality. In this way, the delivery of a higher amount of the tetraoxane moiety, the actual pharmacophore, to the site of action of antimalarial peroxides, the infected erythrocyte,<sup>11</sup> could be realized.

Here, preliminary results on the synthesis of dendrimer cores based on steroidal branches are reported

### SYNTHESIS

The first step in the synthesis is the coupling reaction of trimesoyl trichloride (**1**)<sup>12</sup> (an initial core) with the building block derived from deoxycholic acid. The direct esterification of trimesoyl trichloride by steroidal alcohols, methyl deoxycholate (**2**) and methyl deoxycholate 12 $\alpha$ -acetate (**3**)<sup>13</sup>, in dry toluene and in the presence of DMAP, afforded triester cores **4** and **5** in 56 % and 89 % yield, respectively (Scheme 1).<sup>14</sup> The subsequent step, the selective hydrolysis of the ester at C(24) of the steroidal units using various reac-



Scheme 1.

tion conditions, NaOH/MeOH/H<sub>2</sub>O, LiOH/MeOH/H<sub>2</sub>O, KO<sup>t</sup>Bu/DMSO, NaCN/HMPT, NaCN/Py, gave unsatisfactory results. In addition, demethylation of the methyl ester **5** was effected using the system AlBr<sub>3</sub>/EtSH,<sup>15</sup> yielding the triacid core **6** in only 31 % yield. Hence, a new synthon had to be derived, which would be properly designed for the next coupling reaction and at the same time, affording a product suitably protected for further synthetic manipulations.

Because of their ease of preparation and mild deprotection route, it was decided to use benzyl esters in the further synthesis. Direct esterification of trimesoyl trichloride by the benzyl deoxycholate **7** and benzyl deoxycholate 12 $\alpha$ -benzoate (**8**)<sup>16</sup> afforded the triester cores **9** and **10**, 79 % and 93 % yield, respectively. Heterogeneous hydrogenolysis of compound **10** under mild reaction conditions using the ammonium formate/methanol – Pd/C<sup>17</sup> system, afforded the triacid core **11** in 90 % yield. Reduction of compound **11** by NaBH<sub>4</sub>/BF<sub>3</sub>·Et<sub>2</sub>O gave the triol **12** in 99 % yield.<sup>18</sup> For characterisation purposes, the triol core **12** was acylated to give the triacetate **13**. The 12 $\alpha$ -acetate **14** was prepared from core **9**, using TMSOTf catalysed acylation (79 % yield).

The introduction of the peroxide moiety as well as the results of biological screening will be published at a later time.

## EXPERIMENTAL

### General

Melting points were determined on a Boetius PMHK apparatus and are not corrected. IR spectra were recorded on a Perkin-Elmer Spectrophotometer FT-IR 1725X. <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra were recorded on Varian Gemini-200 spectrometer (at 200 and 50 MHz, respectively) in the indicated solvent using TMS as the internal standard. Chemical shifts are expressed in ppm ( $\delta$ ) values and coupling constants ( $J$ ) in Hz. ESI MS spectra were recorded on an Autospec TOF instrument in the positive ion mode using CH<sub>3</sub>CN/H<sub>2</sub>O (1/1) with 1 % AcOH as the carrying solvent solution. The stock solutions of the samples were diluted in order to obtain 10 pmol/L solutions using the carrying solvent. The source temperature was 75 °C, and the cone voltage was set to 40 V. Abbreviations : Trm – trimesoyl; Bn – benzyl; Bz – benzoyl.

*General procedure for the synthesis of the core.* To a solution of steroidal alcohol (3.3 mmol) and DMAP (9.9 mmol) in dry toluene (60 mL), trimesoyl trichloride (1.00 mmol) was added and heated to reflux until the end of the reaction (TLC: toluene/ethyl acetate 9/1). The reaction mixture was allowed to cool to room temperature and purified by dry-flash chromatography, using the same solvent mixture as eluent.

#### 1,3,5-Tris[(methyl 12 $\alpha$ -hydroxycholanoate)-3 $\alpha$ -oxycarbonyl]benzene (**4**)

Colourless, glassy substance, obtained according to the *General procedure* (56 % yield), mp. 163–173 °C (dichloromethane). IR (KBr): 3545 *m*, 3087 *w*, 2945 *s*, 1724 *s*, 1635 *w*, 1448 *m*, 1381 *m*, 1243 *s*, 1195 *m*, 1170 *m*, 986 *m* cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 8.79 (*s*, 3H–C<sub>Trm</sub>), 5.03 (*m*, 3H–C(3)), 4.00 (*s*, 3H–C(12)), 3.67 (*s*, 9H, CH<sub>3</sub>O<sub>2</sub>C(24)), 0.97 (*s*, 9H, H<sub>3</sub>C–C(10)), 0.70 (*s*, 9H, H<sub>3</sub>C–C(13)). <sup>13</sup>C-NMR-DEPT (50 MHz, CDCl<sub>3</sub>): 174.72 (C(24)), 164.61 (Trm–COOSt), 134.26 (C<sub>Trm</sub>–H), 131.71 (C<sub>Trm</sub>–COOSt), 75.67 (CH–O–), 73.01 (CH–O–), 51.44 (CH<sub>3</sub>O–), 48.23 (CH), 47.21 (CH), 46.41 (C<sub>q</sub>), 41.82 (CH), 35.91 (CH), 35.03 (CH), 34.82 (CH<sub>2</sub>), 34.10 (C<sub>q</sub>), 33.60 (CH), 32.10 (CH<sub>2</sub>), 30.97 (CH<sub>2</sub>), 30.85 (CH<sub>2</sub>), 28.68 (CH<sub>2</sub>), 27.37 (CH<sub>2</sub>), 26.88 (CH<sub>2</sub>), 26.55 (CH<sub>2</sub>), 25.93 (CH<sub>2</sub>), 23.56 (CH<sub>2</sub>), 23.04 (CH<sub>3</sub>–C(10)), 17.25 (CH<sub>3</sub>–C(20)), 12.69 (CH<sub>3</sub>–C(13)).

#### 1,3,5-Tris[(methyl 12 $\alpha$ -acetoxcholanoate)-3 $\alpha$ -oxycarbonyl]benzene (**5**)

Colourless, glassy substance, obtained according to the *General procedure* (89 % yield), mp. 164–170 °C. IR (KBr): 3451 *w*, 2951 *s*, 1738 *s*, 1450 *m*, 1379 *m*, 1244 *s* cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz,

CDCl<sub>3</sub>): 8.82 (s, 3H-C<sub>Tmm</sub>), 5.10 (bs, 3H-C(12)), 5.01 (m, 3H-C(3)), 3.67 (s, CH<sub>3</sub>O<sub>2</sub>C(24)), 2.13 (s, CH<sub>3</sub>COO<sup>-</sup>), 0.96 (s, 9H, H<sub>3</sub>C-C(10)), 0.82 (d, *J* = 5.6 Hz, 9H, H<sub>3</sub>C-C(20)), 0.75 (s, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR-DEPT (50 MHz, CDCl<sub>3</sub>): 174.60 (C(24)), 170.58 (CH<sub>3</sub>COO<sup>-</sup>), 164.77 (Tmm-COOS<sub>t</sub>), 134.30 (C<sub>Tmm</sub>-H), 131.73 (C<sub>Tmm</sub>-COOS<sub>t</sub>), 75.80 (CH-O<sup>-</sup>), 51.44 (CH<sub>3</sub>O<sup>-</sup>), 49.34 (CH), 47.50 (CH), 44.98 (C<sub>q</sub>), 41.90 (CH), 35.65 (CH), 34.63 (CH), 34.47 (CH<sub>2</sub>), 34.03 (CH), 32.27 (CH<sub>2</sub>), 30.90 (CH<sub>2</sub>), 30.76 (CH<sub>2</sub>), 27.28 (CH<sub>2</sub>), 26.90 (CH<sub>2</sub>), 25.71 (CH<sub>2</sub>), 23.34 (CH<sub>2</sub>), 23.04 (CH<sub>3</sub>COO<sup>-</sup>), 21.42 (CH<sub>3</sub>-C(10)), 17.43 (CH<sub>3</sub>-C(20)), 12.35 (CH<sub>3</sub>-C(13)). MS-ESI (*m/z*): 1525 (100 %, M<sup>+</sup>+Na), 1541 (35 %, M<sup>+</sup>+K).

*1,3,5-Tris[(12 $\alpha$ -acetoxycholanoic acid)-3 $\alpha$ -oxycarbonyl]benzene (6)*

AlBr<sub>3</sub> (133 mg, 5.00 mmol) was carefully dissolved in EtSH (10 mL) (under argon) and **5** (500 mg, 0.33 mmol) was added, in one portion. The obtained mixture was stirred for 24 h at room temperature and then methanol (10 mL) was added until the precipitate dissolved. The reaction mixture was then treated with dilute HCl (3.6 %, 20 mL), extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 × 15 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. Dry-flash chromatography on SiO<sub>2</sub> afforded 150.5 mg (31 %) of triacid **6**, as a colourless powder. IR (KBr): 3445 w, 2951 s, 2870 m, 1724 s, 1450 m, 1380 m, 1244 s, 1025 m, 987 m, 745 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 8.83 (s, 3H-C<sub>Tmm</sub>), 5.11 (bs, 3H-C(12)), 5.01 (m, 3H-C(3)), 2.14 (s, 9H, CH<sub>3</sub>COO<sup>-</sup>), 0.96 (s, 9H, H<sub>3</sub>C-C(10)), 0.84 (d, *J* = 2 Hz, 9H, H<sub>3</sub>C-C(20)), 0.75 (s, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR (50 MHz, CDCl<sub>3</sub>): 180.05, 170.69, 164.79, 131.71, 129.00, 75.89, 49.36, 47.54, 44.99, 41.88, 35.65, 34.62, 34.03, 32.27, 30.94, 30.54, 27.30, 26.90, 26.60, 25.73, 23.36, 23.05, 21.43, 17.43, 12.37.

*1,3,5-Tris[(benzyl 12 $\alpha$ -hydroxycholanoate)-3 $\alpha$ -oxycarbonyl]benzene (9)*

Colourless, glassy substance, obtained from diol **7** according to the *General procedure* (50 % yield), mp 110–123 °C. IR (KBr): 3529 m, 3034 w, 2942 s, 2868 s, 1725 s, 1451 m, 1381 m, 1242 s, 1158 m, 1094 w, 1024 w, 985 m, cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 8.79 (s, 3H-C<sub>Tmm</sub>), 7.36–7.31 (5H-C<sub>Ph</sub>), 5.12 (s, Ph-CH<sub>2</sub>O<sup>-</sup>), 5.05 (m, 3H-C(3)), 3.98 (s, 3H-C(12)), 0.96 (s, 9H, H<sub>3</sub>C-C(10)), 0.67 (s, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR-DEPT (50 MHz; CDCl<sub>3</sub>): 174.00 (C(24)), 164.54 (Tmm-COOS<sub>t</sub>), 136.01 (C<sub>Ph</sub>-CH<sub>2</sub>O<sup>-</sup>), 134.18 (C<sub>Tmm</sub>-H), 131.64 (C<sub>Tmm</sub>-COOS<sub>t</sub>), 128.47 (C<sub>Ph</sub>-H), 128.16 (C<sub>Ph</sub>-H), 75.62 (CH-O<sup>-</sup>), 72.94 (CH-O<sup>-</sup>), 66.00 (Ph-CH<sub>2</sub>O<sup>-</sup>), 48.16 (CH), 47.16 (CH), 46.35 (C<sub>q</sub>), 41.77 (CH), 35.84 (CH), 34.94 (CH), 34.94 (CH<sub>2</sub>), 34.05 (C<sub>q</sub>), 33.52 (CH), 32.05 (CH<sub>2</sub>), 31.16 (CH<sub>2</sub>), 30.77 (CH<sub>2</sub>), 28.62 (CH<sub>2</sub>), 27.33 (CH<sub>2</sub>), 26.84 (CH<sub>2</sub>), 26.50 (CH<sub>2</sub>), 25.89 (CH<sub>2</sub>), 23.53 (CH<sub>2</sub>), 23.00 (CH<sub>3</sub>-C(10)), 17.19 (CH<sub>3</sub>-C(20)), 12.64 (CH<sub>3</sub>-C(13)). MS-ESI (*m/z*): 1627 (100 %, M<sup>+</sup>+Na), 1643 (38 %, M<sup>+</sup>+K).

*1,3,5-Tris[(benzyl 12 $\alpha$ -acetoxycholanoate)-3 $\alpha$ -oxycarbonyl]benzene (14)*

To a solution of Ac<sub>2</sub>O (0.050 mL) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL), cooled on ice, one drop of TMSOTf was added. Then, a solution of compound **9** (200 mg, 0.125 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added dropwise. The reaction was completed in 10 min (TLC: PhMe/EtOAc 9/1). The reaction mixture was washed with sat. NaHCO<sub>3</sub> several times, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. After drying *in vacuo* 173.2 mg (79 %) of crude core **14** was obtained. Mp 116–125 °C. IR (KBr): 2947 s, 2868 m, 1732 s, 1244 s, 1158 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz; CDCl<sub>3</sub>): 8.83 (s, 3H-C<sub>Tmm</sub>), 7.36 (m, 15H-C<sub>Ph</sub>), 5.11 (bs, 3H-C(12)), 5.10–4.90 (m, 3H-C(3)), 2.11 (s, 9H, CH<sub>3</sub>COO<sup>-</sup>), 0.96 (s, 9H, H<sub>3</sub>C-C(10)), 0.80 (d, *J* = 5.6 Hz, 9H, H<sub>3</sub>C-C(20)), 0.72 (s, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR (50 MHz; CDCl<sub>3</sub>): 173.97, 170.64, 164.79, 136.03, 134.33, 131.71, 128.52, 128.27, 128.20, 75.85, 75.80, 66.09, 49.33, 47.56, 44.98, 41.90, 35.65, 34.60, 34.47, 34.03, 32.27, 31.17, 30.74, 27.26, 26.91, 26.60, 25.71, 23.35, 21.45, 17.41, 12.35. MS-ESI (*m/z*): 1753 (100 %, M<sup>+</sup>+Na), 1769 (28 %, M<sup>+</sup>+K).

*1,3,5-Tris[(benzyl 12 $\alpha$ -benzoyloxycholanoate)-3 $\alpha$ -oxycarbonyl]benzene (10)*

Colourless, glassy substance, obtained from monoalcohol **8** according to the *General procedure* (93 % yield). Mp 119–129 °C. IR (KBr): 3550–3350 m, 2947 s, 2869 m, 1718 s, 1603 w, 1451 m, 1382 w, 1273 s, 1241 s, 1158 m, 1112 m, 1069 w, 1026 w, 986 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz; CDCl<sub>3</sub>): 8.60 (s, 3H-C<sub>Tmm</sub>), 8.05 (d, *J* = 7 Hz, 6H-C<sub>Bz</sub>), 7.4–7.2 (m, 24H), 5.38 (bs, 3H-C(12)), 5.07 (s, Ph-CH<sub>2</sub>O<sup>-</sup>), 4.90

(*m*, 3H-C(3)), 0.98 (*s*, 9H, H<sub>3</sub>C-C(10)), 0.84 (*bs*, 9H, H<sub>3</sub>C-C(20)), 0.80 (*s*, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR-DEPT (50 MHz; CDCl<sub>3</sub>): 173.82 (C(24)), 165.90 (Ph-COO-), 164.52 (Trm-COOST), 135.95 (C<sub>Ph</sub>-CH<sub>2</sub>O-), 134.02 (C<sub>Trm</sub>-H), 132.85 (C<sub>Bz</sub>-H), 131.44 (C<sub>Trm</sub>-COOST), 130.55 (C<sub>Bz</sub>-COO-), 129.25 (C<sub>Bz</sub>-H), 128.93 (C<sub>Bz</sub>-H), 128.42 (C<sub>Ph</sub>-H), 128.12 (C<sub>Ph</sub>-H), 128.07 (C<sub>Ph</sub>-H), 76.20 (CH-O-), 75.53 (CH-O-), 65.93 (Ph-CH<sub>2</sub>O-), 50.07 (CH), 47.76 (CH), 45.36 (C<sub>q</sub>), 41.75 (CH), 35.64 (CH), 34.63 (CH), 34.63 (CH<sub>2</sub>), 33.91 (C<sub>q</sub>), 32.12 (CH<sub>2</sub>), 31.08 (CH<sub>2</sub>), 30.61 (CH<sub>2</sub>), 27.28 (CH<sub>2</sub>), 26.75 (CH<sub>2</sub>), 26.22 (CH<sub>2</sub>), 25.79 (CH<sub>2</sub>), 23.36 (CH<sub>2</sub>), 22.95 (CH<sub>3</sub>-C(10)), 17.32 (CH<sub>3</sub>-C(20)), 12.46 (CH<sub>3</sub>-C(13)). MS-ESI (*m/z*): 1939 (100 %, M<sup>+</sup>+Na), 1955 (26 %, M<sup>+</sup>+K).

*1,3,5-Tris[(12 $\alpha$ -benzoyloxycholanoic acid)-3 $\alpha$ -oxycarbonyl]benzene (11)*

To a suspension of **10** (117 mg, 0.061 mmol), HCOONH<sub>4</sub> (100 mg) and 10 % Pd/C (100 mg) in ethyl acetate (3 mL), at room temperature, methanol (3 mL) was added. The reaction mixture was stirred at room temperature under argon until the end of the reaction (1 h; TLC: PhMe/acetone 1/1). The mixture was filtered through celite and the filtrate was evaporated. The solid was suspended in CH<sub>2</sub>Cl<sub>2</sub> and washed with brine. After drying over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporating, pasting from benzene and drying *in vacuo* 90.0 mg (90 %) of triacid core **11** was isolated. Mp 201–210 °C (benzene). IR (KBr): 3600–3150 w, 2947 m, 2869 m, 1719 s, 1452 m, 1382 w, 1276 s, 1241 s, 1171 w, 1112 m, 986 w, 715 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz; CDCl<sub>3</sub>): 11.96 (*s*, HOOC(24)), 8.24 (*s*, 3H-C<sub>Trm</sub>), 7.93 (*bs*, 6H-C<sub>Bz</sub>), 7.32 (*bs*, 6H-C<sub>Bz</sub>), 7.17 (*bs*, 3H-C<sub>Bz</sub>), 5.29 (*bs*, 3H-C(12)), 4.83–4.78 (*m*, 3H-C(3)), 0.89 (*s*, 9H, H<sub>3</sub>C-C(10)), 0.77 (*s*, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR (50 MHz; CDCl<sub>3</sub>): 174.92, 165.23, 163.56, 133.20, 132.97, 131.08, 130.50, 129.04, 128.77, 75.98, 75.30, 49.80, 47.88, 45.16, 35.35, 34.35, 33.71, 30.89, 25.79, 22.75, 17.49, 12.43. MS-ESI (*m/z*): 1669 (100 %, M<sup>+</sup>+Na), 1685 (42 %, M<sup>+</sup>+K).

*1,3,5-Tris [(12 $\alpha$ -benzoyloxy-24-hydroxycholan)-3 $\alpha$ -oxycarbonyl]benzene (12)*

To a solution of **11** (300 mg, 0.182 mmol) in THF (5 mL), NaBH<sub>4</sub> (300 mg) was added. After cessation of the reaction, 0.3 mL of BF<sub>3</sub>·Et<sub>2</sub>O was added dropwise and the reaction was completed in 30 min (TLC: toluene/acetone = 1/1). The reaction mixture was evaporated to dryness, the resulting solid suspended in CH<sub>2</sub>Cl<sub>2</sub>, and washed with 3.6 % HCl followed by sat. NaHCO<sub>3</sub> and brine. After evaporation of the solvent, drying and pasting from diethyl ether, 290 mg (99 %) of colourless, powdered core **12** was isolated.

IR (KBr): 3445 m, 2948 s, 2868 s, 1719 s, 1603 w, 1451 m, 1382 m, 1315 m, 1274 s, 1242 s, 1113 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz; CDCl<sub>3</sub>): 8.60 (*s*, 3H-C<sub>Trm</sub>), 8.06 (*d*, *J* = 7.0, 6H-C<sub>Bz</sub>), 7.46–7.27 (*m*, 9H-C<sub>Bz</sub>), 5.40 (*bs*, 3H-C(12)), 4.94 (*m*, 3H-C(3)), 3.74 (*m*, 2H-C(24)), 0.98 (*s*, 9H, H<sub>3</sub>C-C(10)), 0.83 (*s*, 9H, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR-DEPT (50 MHz; CDCl<sub>3</sub>): 166.01 (PhCOO-), 164.59 (Trm-COOST), 134.06 (C<sub>Trm</sub>-H), 132.86 (C<sub>Bz</sub>-H), 131.49 (C<sub>Trm</sub>-COOST), 130.66 (C<sub>Bz</sub>-COOST), 129.31 (C<sub>Bz</sub>-H), 128.49 (C<sub>Bz</sub>-H), 76.36 (CH-O-), 75.62 (CH-O-), 63.33 (C(24)), 50.13 (CH), 47.96 (CH), 45.39 (C<sub>q</sub>), 41.84 (CH), 35.71 (CH), 35.00 (CH), 34.71 (CH), 34.71 (CH<sub>2</sub>), 33.98 (C<sub>q</sub>), 32.19 (CH<sub>2</sub>), 31.58 (CH<sub>2</sub>), 29.32 (CH<sub>2</sub>), 27.44 (CH<sub>2</sub>), 26.84 (CH<sub>2</sub>), 26.28 (CH<sub>2</sub>), 25.95 (CH<sub>2</sub>), 23.44 (CH<sub>2</sub>), 23.02 (CH<sub>3</sub>-C(10)), 17.74 (CH<sub>3</sub>-C(20)), 12.53 (CH<sub>3</sub>-C(13)). MS-ESI (*m/z*): 1604 (37 %, M<sup>+</sup>), 1605 (38 %, M<sup>+</sup>+1), 1627 (100%, M<sup>+</sup>+Na), 1643 (12 %, M<sup>+</sup>+K).

*1,3,5-Tris[(12 $\alpha$ -benzoyloxy-24-acetoxycholan)-3 $\alpha$ -oxycarbonyl]benzene (13)*

To a solution of **11** (24.6 mg, 0.015 mmol) in THF (0.5 mL), NaBH<sub>4</sub> (20 mg) and BF<sub>3</sub>·Et<sub>2</sub>O (0.1 mL) was added and reduction was completed using the above procedure. Crude core **12** was dissolved in pyridine (2 mL) and Ac<sub>2</sub>O (1 mL) and stirred at room temperature for 20 h. The reaction mixture was then poured into an ice-HCl mixture and extracted with chloroform (3 × 5 mL). After evaporating, drying and pasting from benzene, 25 mg (94 %) of **13** was isolated. Mp 70–80 °C. IR (KBr): 2924 s, 2855 s, 1718 s, 1603 w, 1453 m, 1370 w, 1315 w, 1274 s, 1242 s, 1175 m, 1112 m, 715 m cm<sup>-1</sup>. <sup>1</sup>H-NMR (200 MHz; CDCl<sub>3</sub>): 8.60 (*s*, 3H-C<sub>Ar</sub>), 8.06 (*d*, *J* = 4.2 Hz, 6H-C<sub>Bz</sub>), 7.5–7.3 (*m*, 9H-C<sub>Bz</sub>), 5.4 (*bs*, 3H-C(12)), 5.0–4.85 (*m*, 3H-C(3)), 3.97 (*m*, 2H-C(24)), 2.01 (*s*, 9H, CH<sub>3</sub>COO-), 0.99 (*s*, 9H, H<sub>3</sub>C-C(10)), 0.83 (*s*, H<sub>3</sub>C-C(13)). <sup>13</sup>C-NMR (50 MHz; CDCl<sub>3</sub>): 171.23, 166.01, 164.63, 134.11, 132.89, 131.53, 130.71, 129.36, 128.52, 76.36, 75.64, 64.89, 50.16,

47.98, 45.43, 41.88, 35.76, 34.89, 34.76, 34.75, 34.01, 32.32, 31.88, 31.70, 29.65, 29.32, 27.44, 26.86, 26.33, 25.99, 25.22, 23.47, 23.05, 22.65, 20.96, 17.68, 12.57.

*Acknowledgement.* This work was supported by the Ministry of Science, Technologies and Development of Serbia (Project No. 1579).

## ИЗВОД

## СИНТЕЗА ДЕНДРИМЕРСКОГ СТЕРОИДНОГ ЈЕЗГРА

ТАТЈАНА КОП<sup>1</sup>, ГАБРИЈЕЛА ПОКСФАЛВИ<sup>2</sup> И БОГДАН А. ШОЛАЈА<sup>1</sup>

<sup>1</sup>Хемијски факултет, Универзитет у Београду, њ. бр. 158, Београд и <sup>2</sup>Centro di spettrometria di massa proteomica e biomolecolare, Istituto di Scienze dell'Alimentazione, Consiglio Nazionale delle Ricerche, Avellino, Italy

У овом раду приказана је синтеза дендримерског стероидног језгра са естарским карбоксилним и хидроксилним завршецима. Описани приступ синтези омогућава даљу, комплекснију функционализацију једноставним манипулацијама.

(Примљено 26. фебруара 2004)

## REFERENCES

1. G. R. Newkome, C. N. Moorefield, F. Vogtle, *Dendritic Molecules: Concepts, Syntheses and Perspectives*; VCH: Weinheim, 1996
2. T. Nagasaki, O. Kimura, M. Ukon, I. Hamachi, S. Shinkai, S. Arimori, *J. Chem. Soc. Perkin Trans. 1* (1994) 75
3. J. Barbera, B. Donnio, R. Gimenez, D. Guillon, M. Marcos, A. Omenat, J. L. Serrano, *J. Mater. Chem.* **11** (2001) 2808
4. W. Devonport, M. R. Bryce, G. J. Marshallsay, A. J. Moore, L. M. Goldenberg, *J. Mater. Chem.* **8** (1998) 1361
5. G. R. Newkome, V. V. Narayanan, L. A. Godinez, *J. Org. Chem.* **65** (2000) 1643
6. S. Sengupta, S. K. Sadhukhan, *Tetrahedron Lett.* **42** (2001) 3659
7. K. L. Wooley, C. Hawker, J. M. J. Fréchet, F. Wudl, G. Srdanov, S. Shi, C. Li, M. Kao, *J. Am. Chem. Soc.* **115** (1993) 9836
8. G. R. Newkome, E. F. He, C. N. Moorefield, *Chem. Rev.* **99** (1999) 1689
9. S. Hecht, J. M. J. Fréchet, *Angew. Chem. Int. Ed. Engl.* **40** (2001) 74
10. a) N. Todorović, M. Stefanović, B. Tinant, J.-P. Declercq, M. Makler, B. Šolaja, *Steroids* **61** (1996) 688. b) D. Opsenica, G. Pocsfalvi, Z. Juranić, B. Tinant, J.-P. Declercq, D. E. Kyle, W. K. Milhous, B. A. Šolaja, *J. Med. Chem.* **43** (2000) 3274. c) B. A. Šolaja, N. Terzić, G. Pocsfalvi, L. Gerena, B. Tinant, D. Opsenica, W. K. Milhous, *J. Med. Chem.* **45** (2002) 3331. d) D. Opsenica, D. E. Kyle, W. K. Milhous, B. A. Šolaja, *J. Serb. Chem. Soc.* **68** (2003) 291. e) D. Opsenica, G. Angelovski, G. Pocsfalvi, Z. Juranić, Ž. Žižak, D. Kyle, W. K. Milhous, B. A. Šolaja, *Bioorganic & Medicinal Chemistry* **11** (2003) 2761
11. a) S. R. Meschnick, *Artemisinin and Its Derivatives*, in *Antimalarial Chemotherapy*, P. J. Rosenthal, Ed., Humana Press Inc., N. J., 2001, p. 19. b) For recent development in this field see: U. Ecksteln-Ludwig, R. J. Webb, I. D. A van Goethem, J. M. East, A. G. Lee, M. Kimura, P. N. O'Neill, P. G. Bray, S. A. Ward, S. Krishna, *Nature* **424** (2003) 957
12. Trimesoyl-trichloride was synthesized according to Bennet and Wain, *J. Chem. Soc.* (1936) 111
13. Methyl deoxycholate (**2**) was prepared by refluxing either sodium deoxycholate or deoxycholic acid with HCl/MeOH. 12 $\alpha$ -Monoacetate **3** was prepared from the corresponding 3 $\alpha$ ,12 $\alpha$ -diacetate according to ref. 10<sup>b</sup>
14. S. Kohmoto, D. Fukui, T. Nagashima, K. Kishikawa, M. Yamamoto, K. Yamada, *Chem. Commun.* (1996) 1869

15. M. Node, K. Nishide, M. Sai, E. Fujita, *Tetrahedron Lett.* **52** (1978) 5211, M. Node, K. Nishide, M. Sai, K. Fuji, E. Fujita, *J. Org. Chem.* **46** (1981) 1991
16. Benzyl ester was synthesized from sodium deoxycholate using benzyl alcohol in the presence of DMF.  $12\alpha$ -Monobenzoate **8** was prepared directly in a TMSOTf catalyzed acylation reaction using mixed anhydride prepared according to the procedure of J. A. Price, D. S. Tarbell, *Org. Synthesis Coll. Vol. IV*, 285
17. M. K. Anwer, A. F. Spatola, *Synthesis* (1980) 929
18. A. G. Anderson, R. G. Anderson, T. S. Fujita, *J. Org. Chem.* **27** (1962) 4535.