

Terpenes and *n*-Alkanes in Needles of *Pinus cembra*Biljana Nikolić<sup>a\*</sup>, Marina Todosijević<sup>b</sup>, Mihajlo Ratknić<sup>a</sup>, Iris Đorđević<sup>c</sup>, Jovana Stanković<sup>d</sup>, Mirjana Cvetković<sup>d</sup>, Petar D. Marin<sup>e</sup>, and Vele Tešević<sup>b</sup><sup>a</sup>Institute of Forestry, Kneza Višeslava 3, 11000 Belgrade, Serbia<sup>b</sup>Faculty of Chemistry, Studentski trg 16, P.O. Box 158, University of Belgrade, 11000 Belgrade, Serbia<sup>c</sup>Faculty of Veterinary Medicine, Bulevar oslobođenja 18, University of Belgrade, 11000 Belgrade, Serbia<sup>d</sup>Institute for Chemistry, Technology and Metallurgy, Njegoševa 12, University of Belgrade, 11000 Belgrade, Serbia<sup>e</sup>Institute of Botany and Botanical Garden "Jevremovac", Studentski trg 16, Faculty of Biology, University of Belgrade, 11000 Belgrade, Serbia

smikitis@gmail.com

Received: May 21<sup>st</sup>, 2018; Accepted: June 8<sup>th</sup>, 2018

Simultaneous hydrodistillation and extraction of *Pinus cembra* needles from Slovakia was done (via Likens Nickerson apparatus) for the first time. In essential oil extracts 55 compounds were identified, comprising 99.6% of the extract. The most abundant were monoterpene hydrocarbons (71.0%). In the terpene profile  $\alpha$ -pinene, limonene/ $\beta$ -phellandrene, germacrene D,  $\beta$ -pinene, and  $\delta$ -cadinene dominated (53.2%, 11.4%, 9.4%, 4.6%, and 4.3%, respectively). Seven new compounds for *P. cembra*, such as methyl daniellate (0.5%), 1,8 cineole (0.2%) and *trans*-cadinina-1(6),4-diene (0.2%), etc. were found. In needle cuticular wax of *P. cembra* the amount of nonacosan-10-ol was 75.8%. *n*-Alkanes ranged from C<sub>20</sub> to C<sub>35</sub> with the most dominant C<sub>31</sub>, C<sub>29</sub> and C<sub>33</sub> (33.4%, 16.9%, and 9.6%, resp.). Differences in terpene profiles between Slovakian and Greece from one side and Romanian and Polish *cembra* pines on the other side could be the consequence of its disjuncted areal in Carpathian Mountains caused by glaciation and survival of species in different ecological niches. Obtained differences in *n*-alkane profiles among our and literature results could be the consequence of different age of trees.

**Keywords:** *Pinus cembra*, Terpenes, *n*-Alkanes, Nonacosan-10-ol.

*Pinus cembra* L., known as *cembra* pine, Swiss stone pine or Arolla pine, is five-needle soft pine which belongs to family Pinaceae, genus *Pinus*, subgenus *Strobus*, section *Quinquefoliae*, subsection *Strobus* (classification of Germandt et al. [1]). It is glacial relict naturally widespread at high altitudes in the European Alps as well as in the Carpathian Mountains [2]. It grows very slow in its natural area, but could live more than 500 years [3]. There are many references about essential oil composition of pines [4–8], even on population level [9–11], etc. However, there are only few reports of Swiss stone pine dealing with composition of terpenes [12, 13], their enantiomers [14] and/or their antioxidant and antimicrobial activities [2]. The most detailed is the report of Lis et al. [15] who examined essential oils from different parts of *P. cembra*.

Cuticular waxes and *n*-alkanes of various pines had already been published [16–19], even on population level [20]. In the case of *P. cembra* cuticle thickness [21], wax composition [22], and *n*-alkanes of young trees [23], have been published.

**Table 1:** Terpene compounds (in %) in the needles of *Pinus cembra*

Entry	RT <sup>a)</sup>	RI <sup>b)</sup>	Compds	Area(in %)	Class
1	3.274	801	Hexanal	0.2	AC
2	5.561	923	Tricyclene	0.1	MH
3	5.654	926	$\alpha$ -Thujene	tr <sup>c)</sup>	MH
4	5.843	934	$\alpha$ -Pinene	53.2	MH
5	6.249	949	Camphene	0.6	MH
6	6.390	953	Thuja-2,4(10)-diene	tr	MH
7	6.931	973	Sabinene	0.1	MH
8	7.041	978	$\beta$ -Pinene	4.6	MH
9	7.460	984	Myrcene	0.8	MH
10	7.900	1007	$\alpha$ -Phellandrene	tr	MH
11	8.094	1010	$\delta$ -3-Carene	tr	MH
12	8.324	1016	$\alpha$ -Terpinene	tr	MH
13	8.580	1023	<i>p</i> -Cymene	tr	MH
14	8.711	1030	Limonene/ $\beta$ -Phellandrene	11.4	MH
15	8.857	1032	1,8-Cineole	0.2	OM
16	9.838	1057	$\gamma$ -Terpinene	tr	MH

17	10.973	1090	Terpinolene	0.2	MH
18	14.669	1175	Terpinen-4-ol	tr	OM
19	15.263	1192	$\alpha$ -Terpineol	0.2	OM
20	17.191	1235	Thymol methyl ether	0.1	BC
21	19.443	1285	Bornyl acetate	0.3	OM
22	21.734	1139	$\delta$ -Elemene	tr	SH
23	22.292	1351	$\alpha$ -Longipinene	0.5	SH
24	23.442	1377	$\alpha$ -Copaene	0.2	SH
25	24.079	1391	$\beta$ -Cubebene	0.2	SH
26	24.171	1392	$\beta$ -Elemene	0.1	SH
27	24.468	1401	$\beta$ -Longipinene	0.1	SH
28	24.720	1407	Longifolene	tr	SH
29	25.338	1421	( <i>E</i> )- $\beta$ -Caryophyllene	0.4	SH
30	25.757	1431	$\beta$ -Copaene	0.2	SH
31	26.169	1441	Aromadendrene	0.3	SH
32	26.495	1449	<i>trans</i> -Muurolo-3,5-diene	tr	SH
33	26.662	1453	<i>cis</i> -Muurolo-3,5-diene	tr	SH
34	26.788	1456	$\alpha$ -Humulene	tr	SH
35	27.203	1465	<i>cis</i> -Muurolo-4(14),5-diene	0.1	SH
36	27.415	1471	n.i. <sup>d)</sup> 1	0.2	-
37	27.655	1477	<i>trans</i> -Cadinina-1(6),4-diene	0.2	SH
38	27.779	1480	$\gamma$ -Muurolo-4(14),5-diene	1.0	SH
39	27.973	1485	Germacrene D	9.4	SH
40	28.191	1489	$\beta$ -Selinene	tr	SH
41	28.437	1495	<i>trans</i> -Muurolo-4(14),5-diene	0.1	SH
42	28.578	1499	n.i. 2	0.4	-
43	28.632	1500	Bicyclogermacrene	2.1	SH
44	28.787	1504	$\alpha$ -Muurolo-4(14),5-diene	0.6	SH
45	29.074	1510	$\delta$ -Amorphene	tr	SH
46	29.351	1517	$\gamma$ -Cadinene	2.3	SH
47	29.744	1526	$\delta$ -Cadinene	4.3	SH
48	30.098	1535	<i>trans</i> -Cadinina-1,4-diene	0.1	SH
49	30.348	1540	$\alpha$ -Cadinene	0.1	SH
50	30.559	1546	n.i. 3	tr	-
51	31.712	1574	n.i. 4	tr	-
52	31.863	1578	Germacrene D-4-ol	1.4	OS
53	31.939	1580	Spathulenol	0.1	OS
54	33.439	1617	1,10-di-epi-Cubenol	tr	OS
55	33.955	1630	1-epi-Cubenol	0.1	OS
56	34.504	1644	epi- $\alpha$ -Murolo-4(14),5-diene	1.3	OS
57	34.614	1647	$\alpha$ -Muurolo-4(14),5-diene	0.4	OS
58	34.983	1657	$\alpha$ -Cadinol	1.3	OS
59	57.223	2337	Methyl daniellate	0.5	OT

<sup>a)</sup>RT – retention time; <sup>b)</sup>RI – retention index; <sup>c)</sup>tr – traces (< 0.1%); <sup>d)</sup>n.i. – nonidentified.

*P. cembra* from Carpathian Mountains has disjuncted areal. Up to know *n*-alkanes were studied from territory of Romania [2] and Poland [15]. The aim of this study is to examine for the first time essential oil composition, amount of nonacosan-10-ol content and *n*-alkane profile of cembran pine from Slovakia.

In essential oil extracts among 59 compounds detected, 55 were identified, comprising together 99.6% of the extract (Tables 1,2). The most abundant were monoterpene hydrocarbons (71.0%). Seven new compounds for *P. cembra*, such as methyl daniellate (0.5%), 1,8 cineole (0.2%) and *trans*-cadina-1(6),4-diene (0.2%), etc. were found. Terpene profile was as following:  $\alpha$ -pinene >>> limonene/ $\beta$ -phellandrene >germacrene D > $\beta$ -pinene =  $\delta$ -cadinene. (53.2%, 11.4%, 9.4%, 4.6%, 4.3%, respectively. Symbols were explained by Petrakis et al. [24].

**Table 2:** Terpene classes (in %) in the needles of *P. cembra*.

Class	Area (in %)
<b>Total monoterpenes</b>	<b>71.7</b>
Monoterpene hydrocarbons (MH)	71.0
Oxygenated Monoterpenes (OM)	0.7
<b>Total sesquiterpenes</b>	<b>27.1</b>
Sesquiterpene hydrocarbons (SH)	22.5
Oxygenated sesquiterpenes (OS)	4.6
<b>Others</b>	<b>0.8</b>
Aliphatic compounds (AC) <sup>a)</sup>	0.2
Benzenoid compounds (BC) <sup>b)</sup>	0.1
Others (OT)	0.5
<b>Unknown</b>	<b>0.4</b>
<b>Total (%)</b>	<b>100.0</b>

<sup>a)</sup> Aliphatic aldehydes and hydrocarbons; <sup>b)</sup> Cyclic hydrocarbons.

It is obvious that chemotype of *P. cembra* from Slovakia (present results) has more abundant germacrene D and  $\beta$ -pinene (at third and fourth place in profile) comparing to same species from Poland (where  $\delta$ -cadinene is more abundant than germacrene D) [15]. Furthermore, *P. cembra* from Romania [2] differed from both of them with abundant  $\alpha$ -cadinene,  $\gamma$ -cadinene and camphene (at third, fourth and fifth place, resp.).

In needle cuticular wax of *P. cembra* the amount of nonacosan-10-ol was 75.8%, which is significantly higher than in the same species from Swiss Alps [20] where is nearly 60% for one-year and nearly 50% for 2–4 year old needles. In presented work *n*-alkanes ranged from C<sub>20</sub> to C<sub>35</sub> with the most dominant C<sub>31</sub>, C<sub>29</sub>, C<sub>30</sub> and C<sub>33</sub> (33.4%, 16.9%, 10.6% and 9.6%, resp.). In nursery conditions [21] *n*-alkanes of *P. cembra* ranged from C<sub>18</sub> to C<sub>33</sub>, where the most dominant were C<sub>31</sub>, C<sub>27</sub>, C<sub>26</sub>, C<sub>29</sub> and C<sub>25</sub> (20.0%, 6.64%, 5.26%, 5.17% and 5.14%, resp.). Interestingly, in its var. *glauca* range of *n*-alkanes is narrower (C<sub>21</sub> to C<sub>31</sub>), and the most dominant was also C<sub>31</sub>, but with significantly lower abundance (4.51%), followed by C<sub>25</sub>, C<sub>24</sub> and C<sub>23</sub> (4.23%, 4.00% and 3.38%, resp.).

Differences in terpene profiles and *n*-alkane composition among Slovakian and Greek [25] from one side and Romanian and Polish cembran pines on the other side could be the consequence of disjuncted area of *P. cembra* in Carpathian Mountains caused by glaciation as well as survival and development of species in ecological niches. Obtained differences in *n*-alkane profiles among our and literature results could be the consequence of different age of trees.

## Experimental

**Plant material:** Twigs with needles from the lowest third of the full tree crown (up to 60 years old) were collected in autumn 2015 from

Slovakia, Mt. Vysoké Tatry, locality Štrbské Pleso (elevation about 1500 m). The collected twigs were stored at –20°C prior to further analyses.

**Extraction of essential oil:** Essential oil was obtained by simultaneous distillation and extraction with dichloromethane via Likens-Nickerson apparatus.

**GC-FID and GC/MS analyses of essential oil:** GC-FID and GC/MS analyses were carried out with an Agilent 7890A apparatus equipped with an 5975C MSD, FID, and a HP-5MSI fused-silica cap. col. (30 m × 0.25 mm × 0.25  $\mu$ m). The oven temperature was programmed linearly rising from 60 to 315°C for 15 min; injector: 250°C; FID detect.: 300°C; carrier gas, He (1.0 mL/min at 210°C), injection vol. 1  $\mu$ L split ratio, 10:1. EI-MS (70 eV), *m/z* range 40–550.

**Identification of essential oil components:** Identification of all compounds in essential oil was match by comparison of their linear retention indices (relative to C<sub>8</sub>–C<sub>36</sub> *n*-alkanes on the HP-5MSI column) and MS spectra with those of authentic standards from NIST11 and homemade MS library data bases.

**Extraction of needle wax:** The total wax of each sample was extracted by immersing 3 g of leaves in 5 mL of hexane for 45 sec. After extraction the solvent was removed under vacuum at 60°C and the remaining wax dissolved in 1.0 mL hexane. These wax samples were stored at –20°C until further analysis.

**GC and GC-MS analyses of needle wax:** Gas chromatography (GC) and gas chromatography–mass spectrometric (GC-MS) analyses were performed using an Agilent 7890A GC equipped with an inert 5975C XL EI/CI mass selective detector (MSD) and flame ionization detector (FID) connected by capillary flow technology 2-way splitter with make-up. A HP-5MS capillary column (30 m × 0.25 mm × 0.25  $\mu$ m) was used. The GC oven temperature was programmed from 60 to 300 °C at a rate of 3 °C min<sup>-1</sup> and held for 15 min. Helium was used as the carrier gas at 16.255 psi (constant pressure mode). An auto-injection system (Agilent 7683B Series Injector) was employed to inject 1  $\mu$ L of sample. The sample was analyzed in the splitless mode. The injector temperature was 300 °C and the detector temperature 300 °C. MS data was acquired in the EI mode with scan range 30–550 *m/z*, source temperature 230 °C, and quadrupole temperature 150 °C; the solvent delay was 3 min.

**Identification of needle wax components:** The components were identified based on their retention indices and comparison with reference spectra (Wiley and NIST databases) as well as by the retention time locking (RTL) method and the RTL Adams database. The retention indices were experimentally determined using the standard method of Van Den Dool and Kratz [26] involving retention times of *n*-alkanes, injected after the sample under the same chromatographic conditions. The relative abundance of the *n*-alkanes was calculated from the signal intensities of the homologues in the GC-FID traces.

**Acknowledgements** – This research was supported by Grants No. 173029, 173021 and 172053 by the Ministry of Education, Science and Technological Development of the Republic of Serbia. Authors are grateful to Ing. Ján Kučera, NLC-ÚHÚL Zvolen, Slovakia, who collected plant material.

## References

- [1] Gernandt DS, Gaeda López G, Ortiz García S, Liston A. (2005) *Phylogeny and classification of Pinus*. *Taxon*, **54**, 1–13.
- [2] Apetrei LC, Spac A, Brebu M, Tuchilus C, Miron A. (2013) *Composition, and antioxidant and antimicrobial activities of the essential oils of a full-grown Pinus cembra L. tree from the Calimani Mountains (Romania)*. *Journal of the Serbian Chemical Society*, **78**, 27–37.
- [3] Vidaković M. (1982) Četinjače. Morfologija i varijabilnost. JAZU i Sveučilišna naklada Liber, Zagreb, 1–710.
- [4] Rezzi S, Bighelli A, Mouillot D, Casanova J. (2001) *Composition and chemical variability of the essential oil of Pinus nigra subsp. laricio from corsica*. *Flavour and Fragrance Journal*, **16**, 379–383.
- [5] Lahlou M. (2003) *Composition and molluscicidal properties of essential oils of five Moroccan Pinaceae*. *Pharmaceutical Biology*, **41**, 207–210.
- [6] Üstün O, Şenol FS, Kürkçüoğlu M, Orhan IE, Kartal M, Baser KHC. (2012) *Investigation on chemical composition, anticholinesterase and antioxidant activities of extracts and essential oils of Turkish Pinus species and pycnogenol*. *Industrial Crops and Products*, **38**, 115–123.
- [7] Nam AM, Casanova J, Tomi F, Bighelli A. (2014) *Composition and chemical variability of Corsican Pinus halepensis cone oil*. *Natural Product Communications*, **9**, 1361–1364.
- [8] Shpatov AV, Popov SA, Salnikova OI, Khokhrina EA, Shmidt EN, Um BH. (2013) *Low-volatile lipophilic compounds of needles, defoliated twigs, and outer bark of Pinus thunbergii*. *Natural Product Communications*, **8**, 1759–1762.
- [9] Bojovic SR, Jurc M, Drazic DM, Pavlovic P, Mitrovic M, Djurdjevic LA, Dodd R, Afzal-Rafii Z, Barbero M. (2005) *Origin identification of Pinus nigra populations in south western Europe using terpene composition variations*. *Trees*, **19**, 531–538.
- [10] Nikolić B, Ristić M, Bojović S, Marin PD. (2007) *Variability of the needle essential oils of Pinus heldreichii from different populations in Montenegro and Serbia*. *Chemistry and Biodiversity*, **4**, 905–916.
- [11] Šarac Z, Bojović S, Nikolić B, Tešević V, Đorđević I, Marin PD. (2013) *Chemotaxonomic significance of the terpene composition in natural populations of Pinus nigra J.F. Arnold from Serbia*. *Chemistry and Biodiversity*, **10**, 1507–1520.
- [12] Dormont L, Roques A, Malosse C. (1997) *Efficiency of spraying moutain pine cones with oleoresin of Swiss stone pine cones to prevent insect attack*. *Journal of Chemical Ecology*, **23**, 2261–2274.
- [13] Dormont L, Roques A, Malosse C. (1998) *Cone and foliage volatiles emitted by Pinus cembra and some related conifer species*. *Phytochemistry*, **49**, 1269–1277.
- [14] Ochocka JR, Asztemborska M, Sybilska D, Langa W. (2002) *Determination of enantiomers of terpenic hydrocarbons in essential oils obtained from species of Pinus and Abies*. *Pharmaceutical Biology*, **40**, 395–399.
- [15] Lis A, Kalinowska A, Krajewska A, Mellor K. (2017) *Chemical composition of the essential oils from different morphological parts of Pinus cembra L.* *Chemistry and Biodiversity*, **14**, e1600345.
- [16] Herbin GA, Robins PA. (1968) *Studies on plant cuticular waxes.III. The leaf wax alkanes and ω-hydroxy acids of some members of the Cupressaceae and Pinaceae*. *Phytochemistry*, **7**, 1325–1337.
- [17] Cape JN, Fowler D. (1981) *Changes in epicuticular wax of Pinus sylvestris exposed to polluted air*. *Silva Fennica*, **15**, 457–458.
- [18] Burkhardt J, Peters K, Crossley A. (1995) *The presence of structural surface waxes on coniferous needles affects the pattern of dry deposition of fine particles*. *Journal of Experimental Botany*, **46**, 823–831.
- [19] Nikolić B, Tešević V, Đorđević I, Jadranin M, Bojović S, Marin PD. (2010) *n-Alkanes in needle waxes of Pinus heldreichii var. pančići*. *Journal of the Serbian Chemical Society*, **75**, 1337–1346.
- [20] Bojović S, Šarac Z, Nikolić B, Tešević V, Todosijević M, Veljić M, Marin PD. (2012) *Composition of n-alkanes in natural populations of Pinus nigra from Serbia – chemotaxonomic implications*. *Chemistry and Biodiversity*, **9**, 2761–2774.
- [21] Anfodillo T, Di Bisceglie DP, Urso T. (2002) *Minimum cuticular conductance and cuticle features of Picea abies and Pinus cembra needles along analtitudinal gradient in the Dolomites (NE Italian Alps)*. *Tree Physiology*, **22**, 479–487.
- [22] Günthard-Goerg MS. (1986) *Epicuticular wax of needles of Pinus cembra, Pinus sylvestris and Picea abies*. *Europaeen Journal of Forest Pathology*, **16**, 400–408.
- [23] Maffei M, Badino S, Boss S. (2004) *Chemotaxonomic significance of leaf wax n-alkanes in the Pinales (Coniferales)*. *Journal of Biological Research*, **1**, 3–19.
- [24] Petrakis PV, Tsitsimpikou C, Tzakou O, Couladis M, Vagias C, Roussis V. (2001) *Needle volatiles from five Pinus species growing in Greece*. *Flavour and Fragrance Journal*, **16**, 249–252.
- [25] Ioannou E, Koutsaviti A, Tzakou O, Roussis V. (2014) *The genus Pinus: a comparative study on the needle essential oil composition of 46 pine species*. *Phytochemistry Review*, **13**, 741–768.
- [26] Van Den Dool H, Kratz PD. (1963) *A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography*. *Journal of Chromatography*, **11**, 463–471.