10th IAPC Meeting

Tenth World Conference on Physico-Chemical Methods in Drug Discovery &

Sixth World Conference on ADMET and DMPK

Book of Abstracts



10th IAPC Meeting

Tenth World Conference on Physico-Chemical Methods in Drug Discovery & Sixth World Conference on ADMET and DMPK Belgrade, Serbia, September 4-6, 2023

Book of Abstracts

Organized by International Association of Physical Chemists & Faculty of Chemistry, University of Belgrade, Serbia

Published by International Association of Physical Chemists E-mail: office@iapchem.org, URL: http://www.iapchem.org

For Publisher **Zoran Mandić**

Editor Tatjana Verbić & Zoran Mandić

Design, page making and computer layout Aleksandar Dekanski

On Line version only

The Scientific and Organizing Committee:

Tatjana Verbić, Conference Chair University of Belgrade, Serbia,

Alex Avdeef ADME Research, New York, USA

Kiyohiko Sugano Ritsumeikan University, Osaka, Japan

> **Kin Tam** University of Macau, Macau

Zoran Mandić University of Zagreb, Croatia

Klara Valko Biomimetic chromatography Ltd. UK

Godefridus J. Peters Amsterdam University Medical Centers, The Netherlands

> Hong Wan WHDeX Consultng AB, Sweden

Local Organizing Committee

Tatjana Verbić, Conference Chair University of Belgrade, Faculty of Chemistry, Serbia

Goran Roglić University of Belgrade, Faculty of Chemistry, Serbia

Ilija Cvijetić University of Belgrade, Faculty of Chemistry, Serbia

Miloš Pešić University of Belgrade, Faculty of Chemistry, Serbia

Olivera Marković University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

Aleksandar Dekanski University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

> Marija Popović Nikolić University of Belgrade, Faculty of Pharmacy, Serbia

Organization of IAPC-10 Meeting is supported by



Ministry of Science, Technological Development and Innovation of the Republic of Serbia

Sponsors and Exhibitors





<u>N</u>enad M. Kostić Fund for Chemical Sciences

























0 14



Revealing the story of an orphan drug: clofazimine speciation and solubilization as a function of pH

Tatjana Ž. Verbić¹, Alex Avdeef², Kin Y. Tam³, Olivera S. Marković⁴, Miloš P. Pešić¹, Igor A. Topalović¹, Dušan Ž. Veljković¹, Mufaddal Kathawala⁵, Abu T. M. Serajuddin⁵

¹University of Belgrade – Faculty of Chemistry, Belgrade, Serbia, ²in-ADME Research, NY, USA, ³Dept. of Pharmaceutical Sciences, Faculty of Health Sciences, University of Macau, China, ⁴University of Belgrade – Institute of Chemistry, Technology and Metallurgy – National Institute of the Republic of Serbia, Belgrade, Serbia, ⁵St. John's University, College of Pharmacy and Health Sciences, NY, USA

Since the introduction of combinatorial chemistry and high-throughput screening in drug discovery in the early 1990s, the solubility of new chemical entities (NCE) decreased drastically while their lipophilicities increased greatly. Characterizing physicochemical properties of lowsoluble molecules can be especially challenging, since such molecules can undergo complicated reactions in aqueous solution, such as forming precipitates or complexes with buffer species or undergoing self-aggregation (dimer, trimer, etc.)^{1,2} or micelle formations^{3,4}. Most drugs are ionizable. Foremost to the rational interpretation of solution behavior of ionizable drugs in a physiologically-relevant pH domain requires an accurate aqueous pK_{a} , determined by a suitable method. In a pH-dependent measurement of a property (e.g. solubility-, lipophilicity-, permeability-pH), when the apparent pK_a value is different from the true aqueous pK_a value, it may be an early clue that nonideal solution behavior may be taking place. In pharmaceutical research, it may seem cost-effective to use calculated pK_a instead of measured values, but paradoxically, such preference can lead to inaccurate rationalization of the pH-dependent behavior of the drug molecule. For simple molecules, calculated values can be useful, but for today's new drugs or for molecules prone to complicated solution behavior, the use of calculated pK_{as} can substantially wrench the interpretation of solution properties. Clofazimine (CFZ), although discovered about 66 years ago, and used therapeutically for nearly 40 years, exhibits some of the properties of relatively recent drug molecules by being extremely water insoluble and having variable pK_a values reported. We have recently combined potentiometric titrations and UV/Vis spectrophotometry in methanol-water cosolvent media, accompanied by DFT calculations, to assess the hypothesis of CFZ free base dimerization⁵. We reasoned that a soluble dimer might form from drug-drug adhesion along the hydrophobic molecular surface. With lessened exposure of the hydrophobic surface to water, the dimer would be more water soluble than the monomeric free base. In saturated solutions, the apparent solubility in alkaline pH would be elevated due to the presence of the dimer. The effect of that would be a lower pK_a and reverse pK_a cosolvent dependence – the behaviour we have noticed in CFZ aqueous solutions. These findings are of paramount importance for understanding of CFZ speciation and the future progress in developing its improved formulations which is the subject of our ongoing studies.

References:

[1] O.S. Marković, N.G. Patel, A.T.M. Serajuddin, A. Avdeef, T.Ž. Verbić, Mol. Pharmaceutics 2022, 19(2), 710–719.
[2] O.S. Marković, M.P. Pešić, A.V. Shah, A.T.M. Serajuddin, T.Ž. Verbić, A. Avdeef, Eur. J. Pharm. Sci. 2019, 133, 264–274.
[3] A. Avdeef, ADMET & DMPK 2014, 2(1), 33–42.
[4] E. Fuguet, X. Subirats, C. Rafols, E. Bosch, A. Avdeef, Mol. Pharmaceutics 2021, 18(6), 2311–2321, [5] T.Ž. Verbić, K.Y. Tam, D.Ž. Veljković, A.T.M. Serajuddin, A. Avdeef, Mol. Pharmaceutics 2023, 20 (6), 3160-3169.

Acknowledgements: This research has been financially supported by the Ministry of Science, Technological Development and Innovation of Republic of Serbia (Contract No: 451-03-47/2023-01/200026, 451-03-47/2023-01/200168).