

**8<sup>th</sup> Conference of Young Chemists of Serbia**  
**Book of Abstracts**

**29<sup>th</sup> October 2022**  
**University of Belgrade, Faculty of Chemistry**

CIP – Kategorizacija u publikaciji  
Народна библиотека Србије, Београд

54(048)  
577.1(048)  
60(048)  
66.017/.018(048)

CONFERENCE of the Young Chemists of Serbia (8 ; 2022 ; Beograd) Book of abstracts / 8<sup>th</sup> Conference of the Young Chemists of Serbia, [Belgrade], 29<sup>th</sup> October 2022; [organized by Serbian Chemical Society [and] Serbian Young Chemists Club]; [editors Tamara Todorović ... [et al.]]. - Belgrade : Serbian Chemical Society : Serbian Young Chemists Club, 2022 (Belgrade : Development and Research Centre of Graphic Engineering Faculty of Technology and Metallurgy). - 150 str. : ilustr. + 24 cm Tiraž 20. - Bibliografija uz većinu apstrakata. - Registar. ISBN 978-86-7132-080-1

1. Srpsko hemijsko društvo (Beograd) 2. Klub mladih hemičara Srbije (Beograd)

a) Хемија - Апстракти b) Биохемија - Апстракти c) Биотехнологија - Апстракти d) Наука о материјалима – Апстракти

COBISS.SR-ID 78648585

***8<sup>th</sup> Conference of Young Chemists of Serbia***

Belgrade, 29<sup>th</sup> October 2022

***Book of Abstracts***

*Published and organized by*

**Serbian Chemical Society and Serbian Young Chemists' Club**

Karnegijeva 4/III, 11000 Belgrade, Serbia

Tel./fax: +381 11 3370 467; [www.shd.org.rs](http://www.shd.org.rs); [office@shd.org.rs](mailto:office@shd.org.rs)

*Publisher*

Dušan **SLADIĆ**, president of Serbian Chemical Society

*Editors*

Jelena **MILOVANOVIĆ**

Marko **RODIĆ**

Vuk **FILIPOVIĆ**

Života **SELAKOVIĆ**

Jelena **KESIĆ**

Mila **LAZOVIĆ**

Mihajlo **JAKANOVSKI**

*Page Layout and Design*

Vuk **FILIPOVIĆ**

Jelena **KESIĆ**

Mila **LAZOVIĆ**

Mihajlo **JAKANOVSKI**

*Circulation*

20 copies

**ISBN 978-86-7132-080-1**

*Printing*

**Development and Research Centre of Graphic Engineering**

Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade, Serbia

## Influence of the presence of halogen substituents on high-energy properties of nitroaromatic molecules

Aleksandra B. Đunović<sup>1</sup>, Ivana S. Veljković<sup>2</sup>, Dušan Ž. Veljković<sup>3</sup>

<sup>1</sup> Innovation Center of the Faculty of Chemistry, Belgrade, Serbia

<sup>2</sup> University of Belgrade, Institute of Chemistry, Technology and Metallurgy, National Institute of the Republic of Serbia, Belgrade, Serbia

<sup>3</sup> University of Belgrade, Faculty of Chemistry, Belgrade, Serbia

Sensitivity towards detonation of high energetic materials (HEMs) and the positive potential in the central regions of their molecular surfaces are directly related. The presence of halogen atoms in HEMs creates the possibility for halogen bonding which can be used for modifying of electrostatic potential values [1]. Also, it has been noticed that the substitution of hydrogen atoms by halogen atoms in molecules like nitromethane leads to a decrease of bond dissociation energy values (BDE) for the C–N bond [2].

In this paper, the geometries and potentials in the central regions of molecular surfaces of 1,4-dihalo-5,8-dinitronaphthalene and 2,3-dihalo-5,8-dinitronaphthalene were analyzed. Optimal geometries and maps of electrostatic potential (MEP) were calculated using PBE/PBE/6-311G\*\* level of theory. The WFA-SAS program was used to obtain MEP for the mentioned molecules. Bond dissociation energies for optimized geometries were calculated using SAPT program.

Results showed that the potentials above the central regions of molecular surfaces in the 2,3-dihalo-5,8-dinitronaphthalene molecules are higher than in the case of 1,4-dihalo-5,8-dinitronaphthalene analogues. The most significant difference was detected in the case of molecules with chlorine as a substituent (up to 3 kcal/mol). However, the dissociation energies of C–N bonds are higher for all 2,3-substituted dinitronaphthalenes compared to 1,4-substituted analogues. There is a decrease in BDE values in both cases, but it is more significant for the 1,4-substituted dinitronaphthalenes, where the BDE value for 1,4-difluoro-5,8-dinitronaphthalene is more than 7 kcal/mol higher compared to the BDE for iodine analogue.

### References

1. A. B. Đunović, D. Ž. Veljković, *CrystEngComm*. **2021**, *23*, 6915.
2. G. M. Khrapkovskii, A. G. Shamov, R. V. Tsyshevsky, D. V. Chachkov, D. L. Egorov, I. V. Aristov, *Comput. Theor. Chem*. **2012**, *985*, 80.

### Acknowledgments

This research was supported by the Science Fund of the Republic of Serbia, PROMIS, #6066886, CD-HEM.