

EMEC21

21st European Meeting on Environmental Chemistry
November 30 – December 3, 2021, Novi Sad, Serbia

www.emec21.rs



Association of Chemistry
and the Environment



Serbian Chemical Society



Matica Srpska

Scientific Committee

Jan Schwarzbauer, president

Organisational Committee


Branimir Jovančičević, president

Executive Committee

Vladimir Beškoski, president



BOOK OF ABSTRACTS



21st European Meeting
on Environmental Chemistry

EMEC 21

Book of Abstracts
21st European Meeting on Environmental Chemistry

Publisher

Serbian Chemical Society
Karnegijeva 4/III, Belgrade, Republic of Serbia

For the publisher

Dušan Sladić
President of the Serbian Chemical Society

Editors

Ivana Ivančev-Tumbas
Vladimir P. Beškoski
Aleksandra Šajnović

Cover page photo

Branko Lučić

Design and prepress

Beoživković, Belgrade

Printed by

RIS Studio, Belgrade

Circulation

150

ISBN

978-86-7132-078-8

Year

2021

Scientific Committee

Jan Schwarzbauer, Germany, president

Ivana Ivančev-Tumbas, Serbia, vice president

Marijana Ačanski, Serbia	Đorđe Jovanović, Serbia
Vladimir P. Beškoski, Serbia	Albert T. Lebedev, Russia
Jelena Bošković, Serbia	Milan Matavulj, Serbia
Anne-Marie Delort, France	Milica Balaban, Bosnia and Herzegovina
Stuart Gibb, UK	Polonca Trebše, Slovenia
Branimir Jovančićević, Serbia	

Organisational Committee

Branimir Jovančićević, president

Maja Turk Sekulić, vice president

Mira Aničić-Urošević	Jovana Orlić
Mališa Antić	Kristijan Pastor
Jelena Avdalović	Ivana Perović
Tanja Brdarić	Petar Pfindt
Aleksandar Đorđević	Srđan Pokorni
Rada Đurović Pejčev	Bojan Radak
Gordana Gajica	Dubravka Relić
Ljubiša Ignjatović	Goran Roglić
Marko Ilić	Sanja Sakan
Konstantin Ilijević	Sladana Savić
Ivana Ivančev-Tumbas	Jelena Savović
Kristina Joksimović	Slavka Stanković
Vladan Joldžić	Vesna Stanković
Milica Kašanin-Grubin	Sanja Stojadinović
Branka Lončarević	Aleksandra Šajnović
Nikoleta Lugonja	Tatjana Šolević Knudsen
Marija Lješević	Anđelka Tomašević
Snežana Maletić	Aleksandra Tubić
Aleksandra J. Mihajlidi-Zelić	Gorica Veselinović
Maja Milanović	Nenad Zarić
Srđan Miletić	Vesna Zlatanović Tomašević
Tijana Miličević	Aleksandra Žerađanin
Dubravka Milovanović	Sanja Živković
Miloš Momčilović	

Executive Committee

Vladimir P. Beškoski, president

Jelena Radonić, vice president

Gordana Gajica

Kristina Joksimović

Milica Kašanin-Grubin

Srđan Miletić

Gorica Veselinović

EMEC21 Secretariat


Aleksandra Šajnović

Aleksandra Tubić

Nevena Antić

The Effects of Microbial Polysaccharides on the Copper Accumulation in *Daphnia magna*

B. Lončarević^{1,*}, M. Lješević¹, M. Marković^{1,2}, G. Gojgić-Cvijović¹, I. Anđelković², V.P. Beškoski³. (1) University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11000, Belgrade, (2) School of Agriculture, Food and Wine, University of Adelaide, Urrbrae SA, 5064, Australia, (3) University of Belgrade, Faculty of Chemistry, Studentski Trg 12-16, Belgrade, 11000, Serbia. *branka.loncarevic@ihtm.bg.ac.rs.



Copper is one of the leading metal pollutants in the water, which can cause adverse effects when present in high concentrations. The *Daphnia magna* is a model organism usually used for the determination of ecotoxicological effects of various compounds since it is highly sensitive to toxic compounds [1].

The aim of this work was to investigate the potential application of microbial extracellular polysaccharides (EPS), levan and pullulan, as agents for reducing the copper toxicity to *D. magna*. The protective effects of EPS were estimated based on the accumulation of copper in the *D. magna* cells.

Levan is a branched fructane EPS [2] and the one used in this study was produced by *Bacillus licheniformis* NS032. Pullulan, a linear glucan EPS [3], was produced by *Aureobasidium pullulans* CH-1. The *D. magna* were exposed to 50 µg/dm³ of Cu (II) or a combination with 50 mg/dm³ and 100 mg/dm³ of levan or

pullulan for 48h in the acute test. Additionally, the prolonged test was performed, where the daphnia were exposed to a 10 µg/dm³ of Cu (II) with or without 50 mg/dm³ of levan or pullulan for 5 days. After the exposure period, the samples were digested and the accumulation of copper in *D. magna* was analysed using the iCAP Qc ICP-MS (Thermo Scientific, United Kingdom).

The results showed that animals exposed to Cu (II) only, accumulated Cu (II) in a greater amount after the prolonged test compared to the acute one, despite the lower concentration. The treatment with EPS during the acute test increased the copper accumulation for both EPS concentrations tested, whereas during the prolonged exposure test, the Cu (II) accumulation was inhibited.

Considering that protective effects of levan and pullulan were observed only with lower copper concentrations and 5 days of exposure, additional experiments are necessary to determine the mechanism of EPS action in order to confirm their possible use as protective agents.

Acknowledgements

This work was supported by the Ministry of Education, Science and Technological Development (MoESTD) of Republic of Serbia (Grants No: 451-03-9/2021-14/200168 and 451-03-9/2021-14/200026).

References

- [1] B. Lončarević, M. Lješević, M. Marković, I. Anđelković, G. Gojgić-Cvijović, D. Jakovljević, V. Beškoski, *Ecotoxicology and Environmental Safety*, 181 (2019) 187-193.
- [2] Á. González-Garcinuño, A. Tabernero, Á. Domínguez, M. A. Galán, E. M. Martín del Valle, *Biotransformation and Biotransformation*, 36 (2018) 233-244.
- [3] K. R. Sugumaran, V. Ponnusami, *Carbohydrate Polymers*, 173 (2017) 573-591.