EMEC21

21st European Meeting on Environmental Chemistry November 30 – December 3, 2021, Novi Sad, Serbia







Matica Srpska

www.emec21.rs



BOOK OF ABSTRACTS



21st European Meeting on Environmental Chemistry

EMEC 21

Book of Abstracts 21st European Meeting on Environmental Chemistry

Publisher

Serbian Chemical Society

Karnegijeva 4/III, Belgrade, Republic of Serbia

For the publisher

Dušan Sladić

President of the Serbian Chemical Society

Editors

Ivana Ivančev-Tumbas

Vladimir P. Beškoski

Aleksandra Šajnović

Cover page photo

Branko Lučić

Design and prepress

Beoživković, Belgrade

Printed by

RIS Studio, Belgrade

Circulation

150

ISBN

978-86-7132-078-8

Year

2021

Scientific Committee

Jan Schwarzbauer, Germany, president Ivana Ivančev-Tumbas, Serbia, vice president

Marijana Ačanski, Serbia Đorđe Jovanović, Serbia Vladimir P. Beškoski, Serbia Albert T. Lebedev, Russia Jelena Bošković, Serbia Milan Matavulj, Serbia

Anne-Marie Delort, France Milica Balaban, Bosnia and Herzegovina

Stuart Gibb, UK Polonca Trebše, Slovenia

Organisational Committee

Branimir Jovančićević, president Maja Turk Sekulić, vice president

Mira Aničić-Urošević Jovana Orlić

Mališa Antić Kristijan Pastor

Jelena Avdalović Ivana Perović

Tanja Brdarić Petar Pfendt

Aleksandar Đorđević Srđan Pokorni
Rada Đurović Pejčev Bojan Radak
Gordana Gajica Dubravka Relić
Ljubiša Ignjatović Goran Roglić

Marko Ilić Sanja Sakan
Konstantin Ilijević Slađana Savić
Ivana Ivančev-Tumbas Jelena Savović

Kristina Joksimović Slavka Stanković
Vladan Joldžić Vesna Stanković
Milica Kašanin-Grubin Sanja Stojadinović
Branka Lončarević Aleksadra Šajnović

Nikoleta Lugonja Tatjana Šolević Knudsen
Marija Lješević Anđelka Tomašević
Snežana Maletić Aleksandra Tubić
Aleksandra J. Mihajlidi-Zelić Gorica Veselinović

Maja Milanović Nenad Zarić

Srđan Miletić Vesna Zlatanović Tomašević
Tijana Milićević Aleksandra Žerađanin
Dubravka Milovanović Sanja Živković

Miloš Momčilović

Branimir Jovančićević, Serbia

Executive Committee

Vladimir P. Beškoski, president Jelena Radonić, vice president

Gordana Gajica Kristina Joksimović Milica Kašanin-Grubin


Srđan Miletić Gorica Veselinović

EMEC21 Secretariat

Aleksandra Šajnović Aleksandra Tubić Nevena Antić

Metabolomic Study of the Biodegradation Pathway of Sodium-Benzoate in *Pseudomonas aeruginosa* san ai

A. Medić^{1,*}, M. Lješević², I. Karadžić¹. (1) University of Belgrade, Faculty of Medicine, Department of Chemistry, Belgrade, Serbia, (2) University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Department of Chemistry, Njegoševa 12, 11000 Belgrade, Serbia; *ana.medic@med.bg.ac.rs.



Many aromatic compounds are considered to be environmental pollutants that can adversely affect flora and fauna, resulting in the entry of toxic compounds into the food chain causing serious health problems and genetic damage in humans.

Fourteen different pathways of aromatic catabolism in *Pseudomonas* have been confirmed [1]. Benzoate is often used as a model compound to investigate the possibility of microbial degradation of aromatic compounds because it is the simplest known aromatic intermediate in the biodegradation of various complex polyaromatic compounds. The obtained information on the bacterial degradation of benzoate can be further used to understand and predict the degradation pathways of more complex aromatic compounds.

Analytical omics methods enable the study of early molecular changes in the organism to sources of pollution and as such can be used to identify a specific metabolic response to a toxic substance, detect new biomarkers and predict the effects of pollutants on organisms and the environment [2].

The goal of this study was to analyze the products of benzoate degradation by polyextremophilic, hydrocarbonoclastic *Pseudomonas aeruginosa* san ai [3] using

targeted metabolomic analysis in order to determine the specificity of the metabolic pathway of sodium-benzoate. The metabolites of benzoate degradation, after 11 h, 24 h, 48 h and 7 days of incubation, were analyzed using a GC x GC-MS approach. The mass spectrum and retention time of compounds clearly confirmed presence of seven benzoate metabolites such as: 3,4-dixydroxybenzoate, catechol, *cis*, *cis*-muconic acid, muconolactone, β -ketoadipate enol-lactone and succinic acid as the end product of the benzoate transformation. Our results indicate the degradation of NaB through the catechol branch of β -ketoadipate degradation pathway, followed with *ortho*- cleavage of catechol. Furthermore, the identified protocatechuate implies the existence of the second branch of β -ketoadipate pathway.

Metabolomic study showed that almost 99% of benzoate was removed / metabolized within 48 hours and clearly indicates that aromatic degradation occurs via β -ketoadipate. *P. aeruginosa* san ai can be considered as a good candidate for application in bioremediation of environments polluted by different aromatic compounds.

Acknowledgements

This study was supported by the Ministry of Education, Science and Technological Development, Republic of Serbia, Projects III 43004 and 176006.

References

[1] J. Nogales, J.L. García, E. Díaz, in *Aerobic Utilization of Hydrocarbons, Oils, and Lipids*, F. Rojo, (Ed.), Springer, Cham, 2017, 1-49.

[2] D. Gouveia, C. Almunia, Y. Cogne, O. Pible, D. Degli-Esposti, A. Salvador, S. Cristobal, D. Sheehan, A. Chaumot, O. Geffard, J. Armengaud, *Journal of Proteomics*, 198 (2019) 66-77.

[3] I. Karadžić, M. Rikalović, L. Izrael-Živković, A. Medić, in *Unique ecosystems - amazing microbes, series extreme environments*, A. Panday, (Ed.), CRC Press (Taylor & Francis group), 2020, 343-358.